

1 Cleaning and pre-treatment procedures for biogenic and synthetic calcium carbonate
2 powders for determination of elemental and boron isotopic compositions

3

4 M. Holcomb^{1,2,3,*}, T.M. DeCarlo⁴, V. Schoepf^{1,2}, D. Dissard^{1,2,5}, K. Tanaka^{1,6,7}, M.
5 McCulloch^{1,2}

6

7 ¹The UWA Oceans Institute and School of Earth and Environment, The University of
8 Western Australia, Crawley 6009, WA, Australia.

9

10 ²ARC Centre of Excellence in Coral Reef Studies, The University of Western Australia,
11 Crawley 6009, WA, Australia.

12

13 ³Centre Scientifique de Monaco, Monaco, Principality of Monaco

14

15 ⁴Massachusetts Institute of Technology/Woods Hole Oceanographic Institution Joint
16 Program in Oceanography/Applied Ocean Physics and Department of Marine Geology
17 and Geophysics, Woods Hole, MA 02543, U.S.A.

18

19 ⁵ Current address: IRD-Sorbonne Universités (UPMC, Univ Paris 06)-CNRS-MNHN,
20 LOCEAN Laboratory, IRD France-Nord, 32, avenue Henri Varagnat, F-93143 Bondy,
21 France

22

⁶Department of Natural History Sciences, Faculty of Science, Hokkaido University, N10
W8, Sapporo 060-0810, Japan.

⁷Current address: Atmosphere and Ocean Research Institute, University of Tokyo,
Kashiwa, Chiba 277-8564, Japan

*Correspondence: mholcomb3051@gmail.com

keywords: coral boron bleach cleaning peroxide aragonite

Abstract

In preparing calcium carbonate samples for the measurement of various geochemical
proxies, it is often necessary to remove contaminating phases while leaving the phase of
interest altered as little as possible. Here we evaluate the effects of some common
cleaning protocols (rinsing (H₂O), bleach (~3% NaOCl), hydrogen peroxide (30%),
sodium hydroxide (0.006 – 0.1 M NaOH), and acid leaching (0.05 N HNO₃)) on the
elemental (Li, B, Na, Mg, Sr, Ba, Pb, and U) and boron isotope composition of both
biogenic and synthetic calcium carbonates formed in marine environments.

In untreated samples, the presence of elevated concentrations of Na and Mg, the most
abundant cations in seawater, can be reduced with minimal cleaning (e.g. rinsing).

Cleaning protocols that cause partial dissolution are problematic, especially for samples
that are compositionally heterogeneous because the remaining sample may be biased
towards particular phases with distinctive elemental or isotopic compositions. We show

that the use of either acid or unbuffered hydrogen peroxide can lead to partial dissolution which was associated with an increase in the U/Ca ratio of the remaining sample. Bleaching or rinsing with water did not result in significant sample dissolution, suggesting these cleaning techniques may be safely used on heterogeneous samples. Cleaning treatments, other than those resulting in significant dissolution of heterogeneous samples, had no significant effect on $\delta^{11}\text{B}$, suggesting that boron isotopes are generally robust to the effects of sample pre-treatment.

1. Introduction

The elemental and isotopic compositions of a wide range of marine calcium carbonates provide insight into mineralization processes (e.g. Elderfield et al., 1996; Allison and Finch, 2010; McCulloch et al., 2012) and are used as proxy records of environmental conditions (e.g. Montagna et al., 2007; LaVigne et al., 2008; Saenger et al., 2008; Zinke et al., 2014). In order to interpret elemental and isotopic compositions, it is critical to ensure that measurements reflect the carbonate material of interest and not a contaminating phase. Marine samples are particularly susceptible to contamination from residual seawater, and biologically deposited materials may contain organic matter which can retain a wide range of elements (Amiel et al., 1973a,b; McConchie and Harriott, 1992; Mohnen 2008; Blanco-Ameijeiras et al., 2012). In some cases, secondary precipitates may form after the initial material of interest was deposited, surfaces may become enriched in certain elements as a result of adsorption, and particles present in the environment may become mixed with or even incorporated into the sample (Barnard et al., 1974; Macintyre et al., 1976; Shen and Boyle 1988; Millero et al., 2001; Nothdurft et

al., 2005). A wide range of cleaning protocols has been developed to remove many of these sources of contamination (Table 1).

Critical to any cleaning method is the need to not only remove contaminating phases, but also to leave the material of interest as pristine as possible. Although many cleaning methods have been extensively tested on samples collected from the natural environment, less attention has been devoted to testing cleaning methods on nominally ‘clean’ materials, (i.e. free from biologically produced organic compounds) to assess the potential for the cleaning protocol to result in the loss of sample or alter compositionally heterogeneous samples. A brief summary of the methods which have been previously tested and the parameters measured is given in Table S1. Unfortunately no cleaning technique is universally applicable for all trace element measurements, and very few studies have addressed the effects of cleaning on $\delta^{11}\text{B}$ (Table S1).

In this study, we used synthetic aragonite precipitated from filtered seawater as well as biogenic samples. For synthetic materials seawater is the most likely contaminant, while seawater and residual organic materials are likely sources of contamination in biogenic samples. Unique to our study is the use of compositionally zoned synthetic aragonite powders, which avoids the complication posed by tissue contamination. This provides an opportunity to assess whether organic removal steps affect aragonite independent of effects from tissue residues. We also give special attention to the effects of cleaning protocols on boron isotopes in both ‘clean’ and ‘organic-rich’ samples, an issue that has to date received relatively little attention.

92 Table 1. Potential contaminants encountered in marine calcium carbonate samples,
 93 methods/reagents for their removal, and example references for each. See Table S1 for a
 94 summary of studies which have compared different cleaning approaches.

Contaminant	Cleaning method	References
Particulate	Manual removal	Barker et al., 2003; van de Flierdt et al., 2010
	Suspension in water/methanol	Boyle and Keigwin, 1985; Barker et al., 2003
Oxides	Reductive cleaning - $\text{NH}_2\text{OH}\cdot\text{HCl}$ and NH_4OH	Boyle, 1981; Blanco-Ameijeiras et al., 2012
	Reductive cleaning - hydrazine and ammonium citrate	Boyle and Keigwin, 1985; Shen and Boyle, 1988
Barite	chelation with alkaline diethylene-triaminepentaacetic acid (DTPA)	Lea and Boyle, 1993
Adsorbed ions	Rinsing	Amiel et al., 1973; Mitsuguchi and Kawakami, 2012
	Weak acid	Shen and Boyle, 1988; Lea and Boyle, 1993; Rosenthal et al., 1997
	Buffers – acetate	Amiel et al., 1973; Tessier et al., 1979; Wang et al., 2013b
	Chelating agents – EDTA	van de Flierdt et al., 2010
Organic	Oxidation with H_2O_2	Sinclair, 2005; Armid et al., 2011; Mitsuguchi and Kawakami, 2012
	Oxidation with H_2O_2 and NaOH	Boyle and Keigwin, 1985; Tudhope et al., 1996; Blanco-Ameijeiras et al., 2012
	Oxidation with H_2O_2 and HClO_4	Cheng et al., 2000
	Oxidation with H_2O_2 and NH_4OH	Foster et al., 2013
	Oxidation with H_2O_2 and $(\text{NaPO}_3)_6$	Hart et al., 1997
	Oxidation with H_2O_2 and $\text{Ca}(\text{OCl})_2$	Fang and Chou, 1992
	Oxidation with H_2O_2 and NaOCl	Bairbakhish et al., 1999
	Oxidation with NaOCl	St. John, 1974; Sanyal et al., 1996; Tudhope et al., 1996; Clode and Marshall, 2003
	Heat (with or without O_2)	Fabry and Delaney, 1989; Abu-Hilal, 1994; Robinson et al., 2004; Wierzbowski, 2007
	Hydrolysis - NaOH	Young, 1971; Pingitore et al., 1993; Clode and Marshall, 2003

	NaClO ₃	Allison et al., 2010
	TMAH (tetramethylammonium hydroxide)	Stoll et al., 2001

2. Methods

2.1. Carbonate samples

Cleaning protocols were tested on a variety of calcium carbonate samples (Table 3).

Samples included relatively clean coral powders derived from *Porites* sp.: NEP, an in-house coral standard derived from a coral core collected from New Ireland (2.5°S, 150.5°E), Papua New Guinea (Alibert and Kinsley 2008), and the inter-laboratory coral standard JCp-1 (Okai et al., 2002). Organic-rich samples included a specimen of *Marginopora vertebralis* (a high Mg-calcite foraminifera) cultured at the University of South Pacific (Fiji) and a sample taken from the tissue zone of a massive *Porites* sp. coral collected from Myrmidon Reef (18.3°S, 147.4°E), Great Barrier Reef, Australia. Both organic-rich samples were dried following collection without any treatment to remove the tissue, thus organic material from the organism remained associated with the calcium carbonate. To test the effects of sample dissolution, compositionally heterogeneous synthetic aragonite samples grown at Woods Hole Oceanographic Institution (WHOI) (see section 2.1.1) were used. A wide range of other biogenic and synthetic aragonite powders from various sources were also used to test the effects of specific cleaning steps on elemental ratios. Samples not already in the form of a powder were ground into a powder using an agate mortar prior to being used for cleaning tests.

2.1.1. Synthetic aragonite

Two synthetic aragonite samples were used for the bulk of the protocol tests. These were prepared by dissolving 3.67 g CaCO_3 , 0.0484 g SrCO_3 , and 17.91 g MgCl_2 in 4.105 kg filtered seawater at ~ 1 atm (~ 101 kPa) pCO_2 . The final solution was passed through a $0.45\ \mu\text{m}$ filter to remove any undissolved particles. A mass of 656.8 ± 0.7 g of filtered solution was added to ~ 1 L PET containers (SOLO) held within a constant temperature bath (25.15°C). Solutions were stirred continuously at 130 rpm with a magnetic stir bar and bubbled with air at either 5 (precipitate 1/19/11 e) or 10 (precipitate 1/19/11 d) ml/min (see supplemental material for further details). Shortly after the start of precipitation (as inferred from initial declines in pH and alkalinity – see Fig. S1), a ^{87}Sr spike was added (per Holcomb et al., 2013) to more than double the ^{87}Sr concentration, while near the end of the experiment a ^{86}Sr spike was added to the solution. Experiments were terminated by collecting the precipitate on a $0.7\ \mu\text{m}$ glass fiber filter, rinsing with distilled water and ethanol, and then air drying the precipitate. The aragonitic composition of the precipitate was confirmed with raman microscopy (per Holcomb et al., 2009).

Additional synthetic aragonites prepared using a range of protocols (e.g. Kitano et al., 1978; Holcomb et al., 2009; Gabitov et al., 2011; Wang et al., 2013a) were also used to test cleaning protocols, all having been rinsed with H_2O and ethanol at the time of collection. Due to the nature of the experiments, all the synthetic aragonites used are expected to be compositionally heterogeneous (e.g. Gaetani and Cohen, 2006; Holcomb et al., 2009), and for experiments spiked with isotope spikes, there are specific Sr isotope heterogeneities corresponding to known time points in the precipitation experiments.

2.2. Cleaning protocols

From the range of existing cleaning methods, we chose five principal cleaning methods to test: rinsing (H_2O), acid washing (HNO_3), bleaching (sodium hypochlorite, NaOCl or equivalently NaClO), oxidation with hydrogen peroxide (H_2O_2), and hydrolysis (NaOH). Even for these five reagents, there are multiple different implementations which have been used in the literature, covering a wide range of reagent strengths, pHs, durations, and temperatures, only a small subset of which were chosen for testing in the current study. Published protocols include H_2O treatments ranging from brief rinses to more than a day (e.g. Allison 1996; Grottoli et al., 2005; Henehan et al., 2013), bleach treatments from ~3 -12% NaOCl (e.g. Allison 1996; Leder et al., 1996; Clode and Marshall 2003; Ingalls et al., 2003), treatment with acids from <0.001 - 0.2 N (e.g. Rosenheim et al., 2004; Brock et al., 2010; Henehan et al., 2013), H_2O_2 treatments at concentrations from <1 - >30% (McConchie and Harriott 1992; Grottoli et al., 2005; Reynaud et al., 2007; Armid et al., 2011; Henehan et al., 2013), and with NaOH <0.1 - ~6 N (e.g. Watanabe et al., 2001; Clode and Marshall 2003; Tambutte et al., 2007; Krause-Nehring et al., 2011). We chose treatments from within the range of published concentrations and durations for testing. A sub-set of the experiments were conducted at the University of Western Australia and the rest at Woods Hole Oceanographic Institution. The different protocols used and samples tested are summarized in Tables 2 and 3.

Table 2. Steps for each cleaning protocol tested. General information relating to multiple protocols is given in section 2.2.1.

Protocol	Steps
----------	-------

1 (bleach)	2-4x H ₂ O rinse, 1x 1-2 mL of 3-7% NaOCl for 15 min including 5 min of sonication, 5-6x H ₂ O rinses including a few seconds of sonication
2 (acid)	protocol 1 followed by 0.5 mL of 0.05 N HNO ₃ for 15 min including 5 min of sonication, 5x H ₂ O rinses
3 (H ₂ O ₂)	2-4x H ₂ O rinse, 1x 1-2 mL of 30% H ₂ O ₂ for 15 min including 5 min of sonication, 5x H ₂ O rinses including a few seconds of sonication
4 (rinsing)	2-4x ~4 mL H ₂ O rinse
5 (sonication)	1 mL H ₂ O for 15 min including 5 min sonication, 4x H ₂ O rinse
6 (bleach x2)	2x H ₂ O rinse, 2x 1-2 mL of 3-7% NaOCl for 30 min including 15 min of sonication, 5x H ₂ O rinses including a few seconds of sonication
7 (NaOH)	2x H ₂ O rinse, 1x 2 mL 0.03 – 0.07 M NaOH for 15 min including 5 min sonication, 5x H ₂ O rinses including a few seconds of sonication
8 (HNO ₃ + bleach)	2x H ₂ O rinse, 1x 0.5 mL of 0.05 N HNO ₃ for 15 min including 5 min of sonication, followed by protocol 1
9 (hot acid)	dissolution in a few drops of ~14 N HNO ₃ , incubated 2 d on a 90 °C hotplate
10 (rinse)	3x H ₂ O rinse
11 (NaOH)	2x 0.02 M NaOH, 2x H ₂ O rinse
12 (NaOH)	3x 0.006 – 0.1 M NaOH, 3x H ₂ O rinse
13 (bleach)	3x 3% NaOCl, 3x H ₂ O rinse

Table 3. Samples analyzed, the minimum number of replicate samples used per protocol, cleaning protocols compared (as detailed in Table 2, sequential protocols are indicated by ‘+’), elements/isotopes measured, laboratory carrying out the analysis, and where the results are shown.

Sample, # replicates	Cleaning protocols	Measurements	Lab	Results
coral (NEP), 3x	none, 1, 2	$\delta^{11}\text{B}$, Li, B, Na, Mg, Sr, Ba, U	UWA	Fig. 1, 2
coral w organics (massive <i>Porites</i>), 1x	none, 1, 4, 6, 9	$\delta^{11}\text{B}$, Mg, Sr, Ba, U	UWA	Fig. 1, S3, section 3.2.4
corals (various), 4x	1, 1+7, 8	$\delta^{11}\text{B}$, Li, B, Na, Mg, Sr, Ba, Pb, U	UWA	Fig. 5, section 3.2.3
coral (JCp-1), 14x	none, 1	$\delta^{11}\text{B}$	UWA	Fig. 1
foraminifera w organics, 2x	1, 3, 4	$\delta^{11}\text{B}$, Li, B, Na, Mg, Sr, Ba, U	UWA	Fig. 1, S2
synthetic aragonite 1/19/11 d, 3x	none, 1, 3, 4, 5	$\delta^{11}\text{B}$, Li, B, Na, Mg, Sr, Ba, Pb, U	UWA	Fig. 1, 4
synthetic aragonite 1/19/11 e, 3x	none, 1, 2	$\delta^{11}\text{B}$, Li, B, Na, Mg, Sr, Ba, Pb, U	UWA	Fig. 1, 3
synthetic aragonite TS3, 3x	1, 4, 7	B, Na, Mg, Ca, Sr, Ba, Pb, U	UWA	Fig. S4
synthetic aragonites (various), 26x	10, 11	Mg, Sr, Ba, U	WHOI	section 3.2.2
synthetic aragonite TS3, 3x	10 x3, 10+12, 11, 10+13	Mg, Sr, Ba, U	WHOI	Fig. S5

2.2.1. Cleaning steps

All treatments carried out at UWA (protocols 1-9) were carried out in a clean room (McCulloch et al., 2014). Sub-samples (~5-20 mg depending on the sample type) of each of the powders were weighed in pre-cleaned, pre-weighed polypropylene tubes (5 mL capacity). At least two sub-samples of each sample type were assigned to a given treatment, except for treatment 9 (Table 3). In addition to samples, empty tubes were also processed as if they were samples to check procedural blanks. During each step of the protocol samples were shaken vigorously during exposure to the given cleaning

agent. All cleaning steps were carried out at room temperature (~21 °C), unless specified otherwise. To separate the powder from the cleaning agent, tubes were centrifuged at ~3000 rcf for 30 – 120 s, depending on the sample, prior to removing the supernatant. Sonication steps were carried out in an Unisonics ultrasonic bath. Water (18 MΩ) came from a Milli-Q Integral 5 unit fitted with progaard TS2 and quantum cartridges (Millipore). Reagents used included ~6 - 14% NaOCl solutions (Merck 105614 or Rowe CS12345), 30% H₂O₂ solution (Rowe CH2852 CC070711), NaOH (Rowe CS12613), HNO₃ was distilled from analytical grade HNO₃ using a Teflon subboiling distillation system (Savillex DST-1000). Once cleaning was completed, tubes were lightly capped and samples dried by placing them on a 60-70°C hotplate with a heat-lamp above the hotplate to keep the tops warm. Tubes were weighed periodically until the weight stabilized, indicating that the sample was dry.

Treatments carried out at WHOI (Table 3) used 0.75 mL volumes of wash/rinse solutions and ~100 µg powder. All processing was carried out in polypropylene 1.5 mL microcentrifuge tubes. Samples were suspended, centrifuged at 10,000 rpm for 2 minutes, and the supernatant pipetted off. Post cleaning, samples were dried in open microcentrifuge tubes placed in a laminar flow hood. Water used for rinsing and preparing reagents was deionized (18 MΩ, Millipore), HNO₃ was trace metal grade (Fisher A509-p212, lot# 1112110), NaOH was from Fisher (975017), and NaOCl was laboratory grade (Fisher ss290).

2.3. Measurements

2.3.1. UWA

Samples were dissolved in HNO₃ (~0.08 N final acid strength) by adding ~0.5 N HNO₃ directly to the sample (in the same tube used for cleaning steps). Solutions were used directly for boron purification (section 2.3.1.1) or diluted in 2% HNO₃ spiked with ~19 ppb Sc, 19 ppb Y, 0.19 ppb Pr, and 0.095 ppb Bi (note 19 ppb V was also included initially, however its use was discontinued in favor of just using Sc for correcting Ca intensities) for trace element measurements. Most trace element measurements were made on ~10 ppm (~0.25 mM) Ca solutions, however to improve measurement precision of ⁷Li, ¹¹B, and ²⁰⁸Pb, some samples were also measured at ~100 ppm (~2.5 mM) Ca – details of which samples were measured at which concentration are given in the supplementary data table.

Trace element measurements were made using a Q-ICP-MS (X-series II, ThermoFisher Scientific) using the standard Xt interface and plasma screen. During measurement, samples were placed in an autosampler (Elemental Scientific SC4 DX). Sample was introduced by a teflon microflow nebulizer (Elemental Scientific PFA-ST) using a 100 µL/min capillary to a quartz spray chamber chilled to 2°C (Elemental Scientific PC3). The nickel cones of the mass spectrometer were preconditioned for a minimum of 30 minutes with either a 10 or 100 ppm Ca (matching the sample concentration to be measured) solution in 2% HNO₃ prior to starting measurements. Analyses were carried out in peak-hopping pulse counting mode. Wash-out was 20 s and uptake 100 s per sample; 2% HNO₃ was used as the wash solution between samples. Measurements used

100 sweeps with 3-10 repetitions (blank measurements were made using fewer
 repetitions than samples or standards) for all measured elements.

Synthetic standards for trace element analysis were prepared from HPS standard
 solutions and used to check measurement reproducibility. A laboratory coral standard
 (*Porites* sp. collected from Davies Reef, Great Barrier Reef, Australia (147.7 °E, 18.8
 °S)) and JCp-1 (Okai et al., 2002) were measured periodically throughout a run to
 standardize measurements. Values for the Davies Reef coral were based on repeated
 measurements against the JCp-1 coral standard. Elemental ratios (mmol/mol) used for
 JCp-1 were as follows: 0.006185 Li/Ca, 0.4596 B/Ca, 19.83 Na/Ca, 4.199 Mg/Ca, 8.838
 Sr/Ca, 0.007465 Ba/Ca, 0.00009106 Pb/Ca, 0.001192 U/Ca based on Okai et al., (2002),
 Inoue et al., (2004, 2009), and Hathorne et al., (2013). For the Davies Reef coral:
 0.00633 Li/Ca, 0.53 B/Ca, 19.8 Na/Ca, 4.674 Mg/Ca, 8.95 Sr/Ca, 0.003965 Ba/Ca,
 0.000126 Pb/Ca, and 0.001096 U/Ca. Measured counts for each element were corrected
 (background subtraction) for background counts (measured on 2% HNO₃ for correcting
 counts of spiked elements, or on the spike solution for correcting other elements).

Expressed as a percentage of typical sample signal intensities, for 10 ppm Ca solutions
 blank values were: 17% ⁷Li, 22% ¹¹B, 0.5% ²³Na, 2% ²⁴ or ²⁵Mg, <0.1% ⁴³Ca, 0.1% ⁴⁵Sc,
 3% ⁸⁴Sr, 0.1% ⁸⁶Sr, <0.1% ⁸⁷Sr, <0.1% ⁸⁸Sr, <0.1% ⁸⁹Y, 2% ¹³⁸Ba, <0.1% ¹⁴¹Pr, 25%
²⁰⁸Pb, 0.1% ²⁰⁹Bi, <0.1% ²³⁸U; and for 100 ppm Ca solutions, blank values were: 0.5%
⁷Li, 5% ¹¹B, 0.2% ²⁵Mg, 0.1% ⁴⁵Sc, 1% ⁴⁶Ca, <0.1% ⁸⁴Sr, <0.1% ⁸⁹Y, 0.1% ¹³⁸Ba, <0.1%
¹⁴¹Pr, 5% ²⁰⁸Pb, 0.1% ²⁰⁹Bi, <0.1% ²³⁸U.

Spikes were used to correct for variations in instrumental mass bias for the measured
 elements closest in mass to a given spike. To calculate sample ratios, count ratios from

two bracketing standards were used, drift between standards was assumed to occur linearly with time. Reproducibility, expressed as percent relative standard deviation (%RSD), for coral standards measured during the analytical runs for which data are presented here was: 4 Li/Ca, 4 B/Ca, 0.5 Na/Ca, 1 Mg/Ca, 0.1-0.4 for Sr/Ca (depending upon the isotope measured), 1 Ba/Ca, and 1 U/Ca.

2.3.1.1 Boron isotopes

Boron was purified using either cation (Bio-Rad AG50W-X8 200-400 mesh) or anion (Bio-Rad AG1-X8 100-200 mesh) exchange resins to adsorb ions from the sample matrix while allowing boric acid to pass through the resin (Holcomb et al., 2014; McCulloch et al., 2014). Measurements were carried out using MC-ICP-MS as described by Holcomb et al. (2014) and McCulloch et al. (2014) using both direct injection and spray chamber introduction systems on either a Neptune Plus (ThermoFisher Scientific) or Nu (Nu Instruments) MC-ICP-MS. Sample measurements were bracketed by SRM-951 or an in-house standard (UWA24.7) in concentration-matched HNO₃ (~0.15 N for most samples). All data are expressed relative to SRM-951. Long-term reproducibility of coral standards was ~0.3‰ (2 standard deviation). For a given sample type, the same extraction and measurement approach was applied to all samples.

2.3.2. WHOI

Samples measured at WHOI were dissolved in 5% HNO₃ to a final concentration of approximately 40 ppm Ca. Counts of ²⁴Mg, ⁴⁸Ca, ⁸⁸Sr, ¹³⁸Ba, and ²³⁸U were measured in low resolution mode on a magnetic sector ICP-MS (Thermo-Finnegan Element2) with 30

scans per sample. Between samples, the sample introduction system was washed for 3 min with 5% HNO₃. Measured counts were background corrected (background subtracted) based on measurements of 5% HNO₃. Counts were converted to elemental ratios based on measurements of JCp-1. Reproducibility, expressed as %RSD, for coral consistency standards was: 1 Mg/Ca, 0.4 Sr/Ca, 0.8 Ba/Ca, and 0.8 U/Ca.


2.4. Statistics

One-way ANOVAs or t-tests were used to detect significant ($p < 0.01$) differences among treatments. Homogeneity of variance assumptions were not always met, however sample sizes were generally similar among treatments. When significant differences were detected with ANOVA, individual treatments were compared using a Scheffe multiple comparison procedure. Test statistics are expressed as t_y or $F_{x,y}$, where x is the treatment degrees of freedom and y is the error degrees of freedom. For treatments where more than ten replicates were made, Grubb's test for outliers was used to identify outliers which were removed from the data prior to further statistical testing. Statistical tests were carried out using the software packages SYSTAT version 9 and SAS version 7.2. Values are reported as mean \pm 2 standard error. Data are provided in a supplemental .xls file.

3. Results:

3.1. Boron isotopes

289 Different cleaning protocols resulted in no significant change in $\delta^{11}\text{B}$ values for most
290 samples (Fig. 1). Only for one synthetic aragonite (1/19/11 e) were there significant
291 differences among treatments (one-way ANOVA $F_{2,6}=14.8$, $p<0.005$), with the acid
292 treated samples having lower $\delta^{11}\text{B}$ than untreated or bleached samples.



294
295 Figure 1. $\delta^{11}\text{B}$ values for different samples (coral powders: NEP, JCP-1, and *Porites* sp.
296 contaminated with tissue, synthetic aragonites (1/19/11 d and 1/19/11 e), and a
297 foraminifera (*M. vertebralis*) contaminated with tissue) prepared with different cleaning
298 protocols. Cleaning protocols used were: untreated (None), rinsed with water (DI),
299 sonicated during water rinse (Soni), NaOCl (Bleach), hydrogen peroxide (H_2O_2), and
300 leaching in 0.05 N HNO_3 (Acid). Synthetic aragonites are plotted against the right hand
301 axis, corals and foraminifera on the left hand axis. Symbols represent means, error bars
302 are 2 standard errors for cleaning replicates; in some instance the error bars are not
303 visible as they are smaller than the symbol.

3.2. Trace elements

3.2.1 'Clean' aragonite

Cleaning the NEP coral powder (Fig. 2) or synthetic aragonite powders (1/19/11 e (Fig. 3), or 1/19/11 d (Fig. 4)) resulted in no significant pair-wise differences in Li/Ca, Ba/Ca, and Sr/Ca (excluding spiked isotopes) ratios. Ratios for $^{86}\text{Sr}/\text{Ca}$ and $^{87}\text{Sr}/\text{Ca}$ differed amongst treatments for synthetic aragonite 1/19/11 d ($F_{4,10}=8.8$, $p<0.005$, and $F_{4,10}=4.8$, $p=0.02$ respectively), with peroxide treated samples tending to have lower $^{86}\text{Sr}/\text{Ca}$ and higher $^{87}\text{Sr}/\text{Ca}$ values than other treatments (Fig. 4). Acid treatment of synthetic aragonite 1/19/11 e also resulted in a significant decline in $^{86}\text{Sr}/\text{Ca}$ ($F_{2,6}=41.3$, $p<0.0005$, pair-wise $p<0.01$, Fig. 3). For the synthetic aragonite treated with peroxide (1/19/11 d), significantly higher Pb/Ca ratios were found following peroxide treatment ($F_{4,10}=37.9$, $p<0.0001$, pair-wise $p<0.0005$). Although not significant, acid treatment of synthetic aragonite 1/19/11 e also tended to increase Pb/Ca ratios. A similar pattern was observed with U/Ca ratios with acid treated samples having significantly higher values than other treatments (ANOVA, $F_{2,5}=200$, $p<0.0001$ for NEP, $F_{2,6}=129$, $p<0.0001$ for 1/19/11 e). For 1/19/11 d U/Ca ratios differed significantly among treatments ($F_{4,10}=14.6$, $p<0.0005$), with bleached samples having significantly lower values than peroxide or sonication treatments ($p<0.005$), as well as tending to be lower than DI washed and untreated ($p<0.05$) samples.

All cleaning protocols led to reductions in Mg/Ca ($F_{2,6}=434$, $p<0.0001$ for NEP, $F_{2,6}=375$, $p<0.0001$ for 1/19/11 e, $F_{4,10}=811$, $p<0.0001$ for 1/19/11 d) and Na/Ca ($F_{2,6}=355$, $p<0.0001$ for NEP, $F_{2,6}=2045$, $p<0.0001$ for 1/19/11 e, $F_{4,10}=506$, $p<0.0001$ for 1/19/11 d) relative to untreated samples. For the synthetic aragonite 1/19/11 d there were

further significant differences amongst treatments: treatment with bleach or hydrogen peroxide resulted in significantly lower Mg/Ca and Na/Ca ratios ($p < 0.005$) than samples that were only rinsed regardless of whether or not samples were sonicated during rinsing. Further samples that were bleached had significantly higher ($p < 0.005$) Mg/Ca ratios than samples treated with peroxide, however Na/Ca values were not significantly higher. For both synthetic aragonite samples (1/19/11 d, e), B/Ca ratios were lower in cleaned ($F_{2,6}=13.0$, $p < 0.01$ for 1/19/11 e, $F_{4,10}=21.7$, $p < 0.0001$ for 1/19/11 d) relative to untreated samples

3.2.2. Importance of rinsing

For all treatments discussed in section 3.2.1, samples were initially rinsed prior to the use of other cleaning agents, primarily to reduce the possibility of soluble ions precipitating out when exposed to cleaning agents, particularly basic cleaning agents. To test whether this was needed, synthetic aragonite samples were either rinsed or exposed directly to NaOH (without prior rinsing). Two outliers (Grubbs outlier test) were removed from the dataset for each of Mg/Ca, Ba/Ca, and Sr/Ca. Relative to samples rinsed with H₂O, samples treated with NaOH without prior rinsing were enriched in Mg/Ca (16% \pm 4, $t_{23}=8.1$, $p < 0.001$), Sr/Ca (4.2% \pm 0.4, $t_{23}=19.1$, $p < 0.001$) and Ba/Ca (18% \pm 3, $t_{23}=11.2$, $p < 0.001$), but depleted in U/Ca (-13% \pm 2, $t_{25}=-10.1$, $p < 0.001$).

3.2.3. Using multiple reagents

Acidic reagents tended to lead to enrichment of elements such as U and Pb (section 3.2.1), whereas basic reagents tended to result in lower values for those same elements


(section 3.2.1, 3.2.2). To test whether enrichment caused by acid treatment could be reversed with subsequent bleaching, several coral samples were either bleached, or treated with acid followed by bleach. The use of an acid wash step prior to bleaching resulted in a slight but significant increase in Na/Ca ($t_{14}=4.9$, $p<0.001$), and a decrease in Ba/Ca ($t_{14}=-4.7$, $p<0.001$), Pb/Ca ($t_{14}=-10.6$, $p<0.001$), and U/Ca ($t_{14}=-4.3$, $p<0.001$) relative to samples exposed to bleach alone. Other ratios were not significantly affected (Fig. 5), nor was $\delta^{11}\text{B}$ (acid washed values were 0.16 ± 0.13 lower $n=9$).

To test whether the effect of basic reagents was cumulative, some samples were either bleached, or bleached and then treated with NaOH. No significant change in isotopic or elemental ratios were detected following NaOH treatment of bleached samples (Fig. 5, $\delta^{11}\text{B}$ reduced by 0.11 ± 0.26 , $n=4$).

3.2.4. Samples with organic matter

In the high-Mg calcite foraminifera containing organic matter, Mg/Ca exhibited modest differences among treatments ($F_{2,3}=13.0$, $p<0.05$), with rinsed samples having slightly higher ($p<0.05$) values than peroxide treated samples (Fig. S2). Na/Ca differed significantly among treatments ($F_{2,3}=49.1$, $p=0.005$), with peroxide treated samples having lower values than either of the other treatments ($p<0.05$). No significant differences among treatments were detected for Sr/Ca, Ba/Ca, Li/Ca, B/Ca, Pb/Ca, or U/Ca.

371



372

373 Figure 2. Element ratios for NEP coral powder prepared with different cleaning
 374 protocols. Values are mmol Sr, Mg, Na, and B, or μmol U, Ba, and Li per mol Ca,
 375 symbols are means, error bars are 2 standard errors.




Figure 3. Element ratios for synthetic aragonite powder (sample 1/19/11 e) prepared with different cleaning protocols. For different Sr isotopes, the Sr/Ca ratio is calculated assuming natural isotopic abundances. Values and symbols are as described for Fig. 2, values for Pb/Ca are $\mu\text{mol}/\text{mol}$.





Figure 4. Element ratios for synthetic aragonite powder (sample 1/19/11 d) prepared with different cleaning protocols. Values are per Fig3.

Significant differences were found among cleaning protocols for Mg/Ca ($F_{3,16}=45.3$, $p<0.001$), Ba/Ca ($F_{3,16}=30.5$, $p<0.001$), and U/Ca ($F_{3,16}=28.7$, $p<0.001$) in the *Porites* sp. sample containing organic matter (Fig. S3). Untreated samples had significantly ($p<0.005$) higher Mg/Ca ratios than samples subject to any of the cleaning procedures and both untreated and rinsed samples had significantly ($p<0.005$) higher Mg/Ca and Ba/Ca ratios than bleached samples regardless of the bleaching duration. Untreated samples exhibited highly variable U/Ca ratios, although U/Ca tended to be lower in untreated samples than bleached samples. Rinsed samples had significantly ($p<0.0001$) lower U/Ca ratios than bleached samples. Treating a rinsed sample with concentrated nitric acid increased U/Ca ratios to values (~ 1.13 $\mu\text{mol/mol}$) comparable to those measured in bleached samples.

3.3. Sample loss

Acid washing led to a significant loss of the sample ($F_{2,6}=29.0$, $p<.001$ for NEP, $F_{2,6}=82.0$, $p<0.0001$ for synthetic aragonite 1/19/11 e, Fig. 6). For the synthetic aragonite powder treated with H_2O_2 (1/19/11 d), the mass was somewhat reduced following treatment ($F_{4,10}=4.9$, $p<0.05$, H_2O_2 lower than untreated $p<.05$). No significant differences were detected among treatments for foraminifera samples.

406




407

408 Figure 5. Mean relative differences (acid + bleach or bleach + NaOH treated
 409 value/bleached value – 1) in M/Ca ratios for different elements measured in various coral
 410 samples. Error bars are 2 standard errors.

411

412



413

414 Figure 6. Relative mass loss (mass lost/sample mass) for different sample types (NEP =
 415 coral, 1/19/11 e = synthetic aragonite, 1/19/11 d = synthetic aragonite, foram =
 416 foraminifera) versus cleaning treatment. Symbols are means, error bars are standard
 417 deviation.

4. Discussion

4.1. Boron isotopes

For the majority of samples tested, no significant differences were found in the $\delta^{11}\text{B}$ values regardless of the cleaning protocol. Only for the synthetic aragonite (1/19/11 e) samples treated with HNO_3 was there a significant decline in $\delta^{11}\text{B}$ (Fig. 1). Dissolution of synthetic aragonite (1/19/11 e) is expected to decrease $\delta^{11}\text{B}$ given the nature of the precipitation experiment. Since aragonite is depleted in $\delta^{11}\text{B}$ relative to the solution from which it forms (e.g. Vengosh et al., 1991; Hemming et al., 1995), the solution becomes enriched in $\delta^{11}\text{B}$ over the course of precipitation. Moreover, the pH increased toward the end of the experiment (Fig. S1), so $\delta^{11}\text{B}$ should increase due to its dependence upon pH (e.g. Klochko et al., 2006). Thus, the final material deposited in the experiment should have the highest $\delta^{11}\text{B}$ values. Based on changes in Sr isotopes (discussed below, Fig. S7), we conclude that the material deposited towards the end of the experiment dissolved to the greatest extent during treatment with HNO_3 .

Seawater contamination would be expected to increase $\delta^{11}\text{B}$ values due to seawater having a $\delta^{11}\text{B}$ of near 40‰. The lack of significant differences between untreated and rinsed samples suggests that seawater contamination did not contribute sufficient boron to affect $\delta^{11}\text{B}$ in the samples tested here. However, in some instances sample treatment reduced B/Ca ratios (section 3.2.1). Since no change in $\delta^{11}\text{B}$ was detected despite changes in B/Ca, this suggests some of the boron removed had an isotopic composition closer to that of the sample than bulk seawater, such as might be expected for adsorbed borate. Residual organic materials present in the coral and

foraminifera samples did not appear to affect $\delta^{11}\text{B}$ values significantly based on the lack of effect of bleach (Fig. 1). For the foraminifera sample, the lack of a significant difference among cleaning treatments may also be due to the relatively low boron concentration in the foraminifera (high Mg-calcite), which is approximately 20% of the concentration measured in a coral (Fig. 2, S2). The relatively low boron concentration led to larger measurement uncertainties, thus limiting our ability to detect any possible effect of cleaning treatments on $\delta^{11}\text{B}$ of foraminifera samples.

Visual assessment following sample dissolution suggested bleached powders had less undissolved residue than either rinsed or peroxide treated samples. However residual organic material had no significant effect on boron isotopes suggesting that relatively little boron with a $\delta^{11}\text{B}$ composition different from the skeleton was contributed by the organic residue. It should be noted that there are organic materials that do contain boron (e.g. Mohnen 2008), so there may be sample types for which residual organics do influence $\delta^{11}\text{B}$, thus it is likely prudent to include an organic removal step as a precautionary measure.

4.2. Trace elements

For all aragonite samples, treatment with any of the cleaning procedures led to a significant reduction in Mg/Ca and Na/Ca, and a slight decline in B/Ca (Fig. 2, 3, 4), consistent with seawater, or weakly adsorbed ions of seawater origin, being a contaminating phase. Sample dissolution is unlikely to be responsible for this pattern as little to no sample dissolution was measured for many of the cleaning procedures (Fig. 6). Samples of the H_2O supernatant post rinsing from the synthetic aragonite 1/19/11 d were

taken and analyzed for elemental composition. The initial rinses had high Na/Ca and Mg/Ca ratios relative to the bulk precipitate (data not shown), suggesting a seawater or weakly adsorbed component with a composition closer to seawater than that of aragonite, consistent with the patterns measured in the precipitates. Since the synthetic aragonites 1/19/11 d and e were spiked with ^{86}Sr and ^{87}Sr at different time points during their growth, sample dissolution should alter the Sr/Ca ratio differently for different isotopes (Fig. S7). Dissolution of the final precipitate to form would reduce ^{86}Sr , since this was the final isotope spike added to the experiments. An increase in ^{87}Sr would suggest dissolution of the initial material formed (nuclei, which tend to have a higher solubility (e.g. Holcomb et al., 2009)), which precipitated prior to addition of the ^{87}Sr spike. In both cases where measurable dissolution of synthetic aragonites 1/19/11 d and e occurred (acid and H_2O_2 treatments), $^{86}\text{Sr}/\text{Ca}$ ratios declined, indicating dissolution of the exterior of the crystals (Fig. S7). Hydrogen peroxide may also attack the site of nucleation, as $^{87}\text{Sr}/\text{Ca}$ tended to increase following peroxide treatment. This could explain, at least in part, why peroxide treated samples tended to have lower Mg/Ca ratios as well (regions of nucleation tend to be enriched in Mg/Ca relative to the surrounding aragonite crystals (e.g. Holcomb et al., 2009)). Such an explanation contrasts with the findings of Mitsuguchi et al. (2001), who observed that peroxide treatment of coral aragonite resulted in higher Na/Ca and Mg/Ca ratios. However, our observation may be related to particle size as the synthetic aragonites used in our study had, based on microscopy (Fig. S6), particle sizes generally less than 15 μm . The nuclei of the synthetic aragonites were relatively more exposed to the cleaning agents than would be the case with larger particle sizes where nuclei could be protected by surrounding fibers, thus allowing preferential

dissolution of the nuclei to occur in the synthetic aragonite. In the work of Mitsuguchi et al. (2001), the typical particle size of crushed coral skeletons was nearly an order of magnitude greater than in our synthetic aragonites. Mitsuguchi et al. (2001) suggested that the large particle size was sufficient to protect centers of calcification from dissolution, and thus lead to enrichment in Na/Ca and Mg/Ca. Acid treatment, however, appears to attack the exterior of the crystal preferentially based on the decline in ^{86}Sr not being associated with a significant change in ^{87}Sr in the synthetic aragonite. Thus, in a more corrosive solution, faster dissolution of the most exposed regions may dominate, leading primarily to a loss of outer crystal edges. For the coral samples treated with acid prior to bleaching, both particle size (not quantified, but individual particles were visible without magnification) and the use of HNO_3 may have contributed to the significant increase in Na/Ca and the trend towards higher Mg/Ca ratios (Fig. 5), consistent with Mitsuguchi et al., (2001). Although particle size represents one potential explanation for differences in patterns between synthetic and biogenic samples, there could also be other differences (e.g. organic compounds) between these samples which contributed to the observed patterns.

Pb/Ca and U/Ca ratios showed similar patterns, tending to be enriched when dissolution occurred, and reduced by treatment with bleach (Figs. 2,3,4). Although the reasons for this behavior are unknown, it may in-part reflect surface adsorption of Pb and U (Fabry and Delaney, 1989; Hong et al., 2002; Godelitsas et al., 2003; Rouff et al., 2005; Du et al., 2011). An enrichment of Pb and U on either the crystal or tube surface may result from exposure to mildly acidic conditions due to sample dissolution releasing U and Pb

from the aragonite - which in-turn may preferentially adsorb onto surfaces, thus leading to enrichment in the remaining sample. Complexation of Pb and U (Turner et al., 1981; Millero et al., 2009) in the bleach solution may help remove ions adsorbed to surfaces. Any enrichment due to acid leaching can potentially be removed by subsequent bleaching, and the combination of acid leaching and bleaching may be more effective than bleach alone at removing any surface enrichment (Fig. 5). However, bleach does not always reduce U/Ca, nor does NaOH (Figs. S4,S5), suggesting that it is neither NaOCl nor high pH that is primarily responsible for removing adsorbed U and Pb. Carbonate (CO_3^{2-}) is known to complex both U and Pb (Turner et al., 1981; Millero et al., 2009) and CO_3^{2-} has been used to remove adsorbed U in other studies (Galun et al., 1983). Carbonate concentrations in the bleach or NaOH solution likely increase over time due to a gradual uptake of atmospheric CO_2 by the solution which converts to CO_3^{2-} under the high pH conditions of the bleach or NaOH solutions. Variations in the CO_3^{2-} concentration of the cleaning solution could thus contribute variability to the change in U/Ca due to bleaching (Figs. S4,S5).

4.3. Organic materials

Although bleaching reduced Ba/Ca ratios in the organic rich coral sample, and longer bleaching treatments resulted in more consistent reductions than shorter treatments (Fig. S3), Ba/Ca remained higher than would generally be expected. This is consistent with other studies which report anomalously high Ba/Ca ratios within the tissue layer of coral cores (e.g. Tudhope et al., 1996; Sinclair et al., 1998; Alibert et al., 2003; Fallon et al., 2003). A wide range of cleaning approaches has been tested to reduce Ba/Ca, however

Ba/Ca ratios generally remain elevated despite cleaning (e.g. Tudhope et al., 1996; Sinclair, 2005). Our observations are consistent with either organic bound Ba (Flor and Moore, 1977; Lea and Boyle, 1993; Reichelte and McOrist, 2003) or a Ba containing phase soluble at high pH (Nagtegaal et al., 2012) making a substantial contribution to the Ba enrichment as bleach greatly reduced the Ba/Ca ratio. However whether the remaining enrichment is due to refractory organic materials or other factors is not known.

The increase in U/Ca following bleaching in organic rich coral samples (Fig. S3) likely reflects adsorption of U by the insoluble organic materials present in the unbleached samples. Though not significant, a similar trend was observed for the foraminifera sample, with both bleach and peroxide treatments being associated with higher U/Ca. Uranium is known to adsorb onto a wide range of organic materials (Amiel et al., 1973b; Galun et al., 1983; Semiao et al., 2010), thus oxidation of those organic materials prior to sample dissolution should increase the concentration of U free in solution. When the organic materials were not oxidized prior to dissolution but rather were oxidized with hot nitric acid post-dissolution, U/Ca ratios were similar (1.13 $\mu\text{mol/mol}$) to those measured in bleached samples (Fig. S3). This suggests adsorption of U by organics as a reason low U/Ca ratios were obtained in samples not treated with an organic oxidation step. For the coral sample used here bleach appears to be sufficient to remove the bulk of the organic material capable of binding U, however for some coral samples bleach alone appears to be insufficient and additional oxidative steps are required to obtain U/Ca values near the range of reported values (data not shown).

4.4. *Cleaning recommendations*

4.4.1. *Rinsing and initial organic removal steps*

Our results confirm the findings of Mitsuguchi et al. (2001) and Watanabe et al. (2001) that sample cleaning treatments help to remove contaminants likely of seawater origin (Mg, Na). Further treatment with bleach reduces the amount of organic material visible in the sample without significant sample dissolution and appears more effective in removing organic materials than H₂O₂ (using similar incubation durations) consistent with many prior comparisons (Pingitore et al., 1993; Gaffey and Bronnimann, 1993; Krause-Nehring et al., 2011; Nagtegaal et al., 2012), though not all (e.g. Pak et al., 2004). For any sample treatment procedure, rinsing with H₂O is a useful first step as it helps to remove residual seawater which might otherwise contribute ions which could form precipitates during subsequent cleaning steps (e.g. section 3.2.2).

Although samples can be rinsed briefly without measureable sample loss (Fig. 6), care should still be taken to avoid excessive exposure to water as both dissolution and recrystallization can occur with prolonged exposure (Amiel et al., 1973a,b; Yoshioka et al., 1986; Pingitore et al., 1993; Holcomb et al., 2009; Shirai et al., 2012). In some instances the pH of the water used for rinsing has been elevated in an effort to reduce dissolution (e.g. Bairbakhish et al., 1999; Blanco-Ameijeiras et al., 2012; Heinmann et al., 2012), however, the effectiveness of this procedure is unknown. The use of sodium hydroxide solutions (~5 M), though effective in reducing dissolution, has been associated with recrystallization (Pingitore et al., 1993; Krause-Nehring et al., 2011), and basic detergent solutions have been found to enhance dissolution (Pingitore et al., 1993), thus elevated pH does not necessarily prevent alteration of the sample during rinsing. It

should be noted that although strong NaOH solutions have been shown to affect samples, much less work has been conducted on how less concentrated bases influence sample composition (Table S1), so weakly basic solutions may satisfactorily reduce dissolution. Treatment with acidic solutions (unbuffered H₂O₂ or HNO₃) can lead to significant dissolution (Pingitore et al., 1993) and alter the composition of heterogeneous samples (Figs. 3,4,6), though there are instances in which some dissolution is considered desirable (e.g. Boiseau and Juillet-Leclerc, 1997). The use of bleach in place of acidic reagents appears advisable for preserving the sample, however for some isotopic ratios ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) bleach may affect values (Grottoli et al., 2005; Wierzbowski, 2007). Other approaches using non-acidic reagents such as buffered H₂O₂ may also provide satisfactory results (e.g. Stoll et al., 2001; Blanco-Ameijeiras et al., 2012). Any cleaning agent should be used with caution as the presence/concentrations of various ions can dramatically affect the rate of ion exchange, dissolution, and recrystallization (Amiel et al., 1973a,b; Yoshioka et al., 1986; Pingitore et al., 1993).

4.4.2. Removing refractory organics

For samples containing refractory organic matter not removed with initial bleach or peroxide treatments an additional oxidation step post-dissolution may be needed for determination of elements such as U. Incubation in hot nitric acid appears to oxidize/hydrolyze sufficient organic material to improve U/Ca values; however this approach was not extensively tested and may not be sufficient for obtaining reliable U/Ca ratios from different sample types. For samples containing refractory organic materials it would be worth testing some of the established oxidation techniques, such as: photo-

oxidation (Spivack and Edmond, 1986; Wara et al., 2003), peroxide treatment (Amiel et al., 1973a; Ihlenfeld et al., 2003), perchloric acid (Tatsumoto and Goldberg, 1959), ashing combined with concentrated HNO_3 and evaporation (Robinson et al., 2004), or concentrated HNO_3 and H_2O_2 (Bohm et al., 2006).

4.4.3. *Removing adsorbed species*

For measurements of species which may adsorb to surfaces (such as Pb and U), we have shown that bleaching can help to remove adsorbed Pb and U, possibly due to carbonate present in the solution. Although carbonate solutions may be useful for removing adsorbed U or Pb species (Galun et al., 1983), there are a number of established protocols for removing these species. Acetate buffers can reduce adsorbed U and have been used for this purpose (Amiel et al., 1973b; Wang et al., 2013b). For Pb and trace metals, short (<2 min) acid wash steps have also been shown to be effective in removing surface contaminants, but longer exposure leads to re-enrichment as the pH of the acid wash solution rises (Shen and Boyle, 1988). As an alternative to extensive sample cleaning procedures, flow through dissolution systems have also been developed for measuring species which tend to adsorb to surfaces (e.g. Klinkhammer et al., 2004).

5. *Conclusions*

For the samples tested, $\delta^{11}\text{B}$ appears to be relatively insensitive to sample cleaning methods, with significant changes occurring only for compositionally heterogeneous samples subjected to dissolution. However other trace elements were more strongly

affected by cleaning. Rinsing samples with H₂O and treatment with NaOCl (bleach) appeared to cause little dissolution. Rinsing helps to remove seawater contamination and/or surface adsorbed Na and Mg, while bleach or H₂O₂ solubilize much of the organic material. These cleaning agents may thus be useful for removing such contaminants from CaCO₃ samples.

Acknowledgements

We wish to thank the following for invaluable assistance with this study: R. Belastock, S. Birdwhistell, E. Bonk, A. Cohen, G. Gaetani, L. Kerr, D. McCorkle, M. Sulanowska, D. Wellwood, and S. White at Woods Hole Oceanographic Institution; D. Allemand, D. Desgré, E. Elia, C. Godinot, N. Segonds, E. Tambutté, S. Tambutté, N. Techer, and A. Venn at the Centre Scientifique de Monaco; J.P. D'Olive Cordero, H. Oskierski, K. Rankenburg, and J. Trotter at the University of Western Australia; and R. Naidu at The University of the South Pacific. Research conducted at the University of Western Australia was supported by the Australian Research Council (ARC) Centre of Excellence for Coral Reef Studies. Research conducted at WHOI was supported by NSF grant OCE-1338320. M.H. was supported by an ARC Super Science Fellowship (at UWA) and a NSF International Research Fellowship (at CSM). T.D. was supported by a NSF Graduate Research Fellowship. M.M. was supported by a Western Australian Premiers Fellowship and an ARC Laureate Fellowship. This study was financially supported in part by Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation (G2301, the Japan Society of the Promotion of Science awarded to KT).

648

649 **References**

- 650 Abu-Hilal, A.H., 1994. Effect of depositional environment and sources of pollution on
651 uranium concentration in sediment, coral, algae and seagrass species from the
652 Gulf of Aqaba (Red Sea). *Marine Pollution Bulletin*, 28: 81-88.
- 653 Alibert, C., Kinsley, L., 2008. A 170-year Sr/Ca and Ba/Ca coral record from the western
654 Pacific warm pool: 1. What can we learn from an unusual coral record? *Journal of*
655 *Geophysical Research*, 113(4): C04008, doi:10.1029/2006JC003979.
- 656 Alibert, C., Kinsley, L., Fallon, S.J., McCulloch, M.T., Berkelmans, R., McAllister, F.,
657 2003. Source of trace element variability in Great Barrier Reef corals affected by
658 the Burdekin flood plumes. *Geochimica et Cosmochimica Acta*, 67(2): 231-246.
- 659 Allison, N., 1996. Geochemical anomalies in coral skeletons and their possible
660 implications for palaeoenvironmental analyses. *Marine Chemistry*, 55: 367-379.
- 661 Allison, N., Finch, A.A., 2010. $\delta^{11}\text{B}$, Sr, Mg and B in a modern *Porites* coral: the
662 relationship between calcification site pH and skeletal chemistry. *Geochimica et*
663 *Cosmochimica Acta*, 74(6): 1790-1800.
- 664 Allison, N., Finch, A.A., EIMF, 2010. The potential origins and palaeoenvironmental
665 implications of high temporal resolution $\delta^{18}\text{O}$ heterogeneity in coral skeletons.
666 *Geochimica et Cosmochimica Acta*, 74: 5537-5548.
- 667 Amiel, A.J., Friedman, G.M., Miller, D.S., 1973a. Distribution and nature of
668 incorporation of trace elements in modern aragonitic corals. *Sedimentology*,
669 20(1): 47-64.

670 Amiel, A.J., Miller, D.S., Friedman, G.M., 1973b. Incorporation of uranium in modern
671 corals. *Sedimentology*, 20(4): 523-528.

672 Armid, A., Asami, R., Fahmiati, T., Sheikh, M.A., Fujimura, H., Higuchi, T., Taira, E.,
673 Shinjo, R., Oomori, T., 2011. Seawater temperature proxies based on D_{Sr} , D_{Mg} ,
674 and D_U from culture experiments using the branching coral *Porites cylindrica*.
675 *Geochimica et Cosmochimica Acta*, 75(15): 4273-4285.

676 Bairbakhish, A.N., Bollmann, J., Sprengel, C., Thierstein, H.R., 1999. Disintegration of
677 aggregates and coccospheres in sediment trap samples. *Marine*
678 *Micropaleontology*, 37(2): 219-223.

679 Barker, S., Greaves, M., Elderfield, H., 2003. A study of cleaning procedures used for
680 foraminiferal Mg/Ca paleothermometry. *Geochemistry, Geophysics, Geosystems*,
681 4(9): 8407.

682 Barnard, L.A., Macintyre, I.G., Pierce, J.W., 1974. Possible environmental index in
683 tropical reef corals. *Nature*, 252: 219-220.

684 Blanco-Ameijeiras, S., Lebrato, M., Stoll, H.M., Iglesias-Rodriguez, M.D., Méndez-
685 Vicente, A., Sett, S., Müller, M.N., Oeschies, A., Schulz, K.G., 2012. Removal of
686 organic magnesium in coccolithophore calcite. *Geochimica et Cosmochimica*
687 *Acta*, 89: 226-239.

688 Bohm, F., Gussone, N., Eisenhauer, A., Dullo, W., Reynaud, S., Paytan, A., 2006.
689 Calcium isotope fractionation in modern scleractinian corals. *Geochimica et*
690 *Cosmochimica Acta*, 70(17): 4452-4462.

691 Boisseau, M., Juillet-Leclerc, A., 1997. H_2O_2 treatment of recent coral aragonite: oxygen
692 and carbon isotopic implications. *Chemical Geology*, 143: 171-180.

693 Boyle, E.A., 1981. Cadmium, zinc, copper, and barium in foraminifera tests. Earth and
694 Planetary Science Letters, 53(1): 11-35.

695 Boyle, E.A., Keigwin, L.D., 1985. Comparison of Atlantic and Pacific paleochemical
696 records for the last 215,000 years: changes in deep ocean circulation and chemical
697 inventories. Earth and Planetary Science Letters, 76(1–2): 135-150.

698 Brock, F., Higham, T., Ditchfield, P., Ramsey, C.B., 2010. Current pretreatment methods
699 for AMS radiocarbon dating at the Oxford radiocarbon accelerator unit (ORAU).
700 Radiocarbon, 52(1): 103-112.

701 Cheng, H., Adkins, J., Edwards, R.L., Boyle, E.A., 2000. U-Th dating of deep-sea corals.
702 Geochimica et Cosmochimica Acta, 64: 2401-2416.

703 Clode, P.L., Marshall, A.T., 2003. Variation in skeletal microstructure of the coral
704 *Galaxea fascicularis*: effects of an aquarium environment and preparatory
705 techniques. The Biological Bulletin, 204: 138-145.

706 Du, Y., Lian, F., Zhu, L., 2011. Biosorption of divalent Pb, Cd and Zn on aragonite and
707 calcite mollusk shells. Environmental Pollution, 159(7): 1763-8.

708 Elderfield, H., Bertram, C.J., Erez, J., 1996. A biomineralization model for the
709 incorporation of trace elements into foraminiferal calcium carbonate. Earth and
710 Planetary Science Letters, 142: 409-423.

711 Fabry, V.J., Delaney, M.L., 1989. Lead-210 and polonium-210 in pteropod and heteropod
712 mollusc shells from the North Pacific: Evaluation of sample treatments and
713 variation with shell size. Journal of Marine Research, 47(4): 933-949.

714 Fallon, S.J., McCulloch, M.T., Alibert, C., 2003. Examining water temperature proxies in
 715 Porites corals from the Great Barrier Reef: a cross-shelf comparison. *Coral Reefs*,
 716 22: 389-404.

717 Fang, L.-S., Chou, Y.-C., 1992. Concentration of fulvic acid in the growth bands of
 718 hermatypic corals in relation to local precipitation. *Coral Reefs*, 11(4): 187-191.

719 Flor, T.H., Moore, W.S., 1977. Radium/Calcium and Uranium/Calcium determinations
 720 for western Atlantic reef corals. *Proceedings of the 3rd International Coral Reef*
 721 *Symposium*: 555-562.

722 Foster, G.L., Hönisch, B., Paris, G., Dwyer, G.S., Rae, J.W.B., Elliott, T., Gaillardet, J.,
 723 Hemming, N.G., Louvat, P., Vengosh, A., 2013. Interlaboratory comparison of
 724 boron isotope analyses of boric acid, seawater and marine CaCO_3 by MC-ICPMS
 725 and NTIMS. *Chemical Geology*, 358: 1-14.

726 Gabitov, R.I., Schmitt, A.K., Rosner, M., McKeegan, K.D., Gaetani, G.A., Cohen, A.L.,
 727 Watson, E.B., Harrison, T.M., 2011. In situ $\delta^7\text{Li}$, Li/Ca, and Mg/Ca analyses of
 728 synthetic aragonites. *Geochemistry Geophysics Geosystems*, 12(3): Q03001,
 729 doi:10.1029/2010GC003322.

730 Gaetani, G.A., Cohen, A.L., 2006. Element partitioning during precipitation of aragonite
 731 from seawater: a framework for understanding paleoproxies. *Geochimica et*
 732 *Cosmochimica Acta*, 70: 4617-4634.

733 Gaffey, S.J., Bronnimann, C.E., 1993. Effects of Bleaching on Organic and Mineral
 734 Phases in Biogenic Carbonates. *Journal of Sedimentary Petrology*, 63: 752-754.

735 Galun, M., Keller, P., Malki, D., Feldstein, H., Galun, E., Siegel, S.M., Siegel, B.Z., 1983.
 736 Removal of Uranium(VI) from Solution by Fungal Biomass and Fungal Wall-
 737 Related Biopolymers. Science (Washington D C), 219: 285-286.

738 Godelitsas, A., Astilleros, J.M., Hallam, K., Harissopoulos, S., Putnis, A., 2003.
 739 Interaction of Calcium Carbonates with Lead in Aqueous Solutions.
 740 Environmental Science & Technology, 37(15): 3351-3360.

741 Grottoli, A.G., Rodrigues, L.J., Matthews, K.A., Palardy, J.E., Gibb, O.T., 2005. Pre-
 742 treatment effects on coral skeletal $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$. Chemical Geology, 221(3-4):
 743 225-242.

744 Hart, S.R., Cohen, A.L., Ramsay, P., 1997. Microscale analysis of Sr/Ca and Ba/Ca in
 745 Porites. Proceedings of the 8th International Coral Reef Symposium, 2: 1707-
 746 1712.

747 Hathorne, E.C., Gagnon, A., Felis, T., Adkins, J., Asami, R., Boer, W., Caillon, N., Case,
 748 D., Cobb, K.M., Douville, E., deMenocal, P., Eisenhauer, A., Garbe-Schönberg,
 749 D., Geibert, W., Goldstein, S., Hughen, K., Inoue, M., Kawahata, H., Kölling, M.,
 750 Cornec, F.L., Linsley, B.K., McGregor, H.V., Montagna, P., Nurhati, I.S., Quinn,
 751 T.M., Raddatz, J., Rebaubier, H., Robinson, L., Sadekov, A., Sherrell, R.,
 752 Sinclair, D., Tudhope, A.W., Wei, G., Wong, H., Wu, H.C., You, C.-F., 2013.
 753 Inter-laboratory study for coral Sr/Ca and other element/Ca ratio measurements.
 754 Geochemistry, Geophysics, Geosystems, 14(9): 3730-3750.

755 Heinemann, A., Fietzke, J., Melzner, F., Böhm, F., Thomsen, J., Garbe-Schönberg, D.,
 756 Eisenhauer, A., 2012. Conditions of *Mytilus edulis* extracellular body fluids and
 757 shell composition in a pH-treatment experiment: Acid-base status, trace elements

758 and $\delta^{11}\text{B}$. *Geochemistry, Geophysics, Geosystems*, 13(1), Q01005,
 759 doi:10.1029/2011GC003790.

760 Hemming, N.G., Reeder, R.J., Hanson, G.N., 1995. Mineral-fluid partitioning and
 761 isotopic fractionation of boron in synthetic calcium carbonate. *Geochimica et*
 762 *Cosmochimica Acta*, 59(2): 371-379.

763 Henehan, M.J., Rae, J.W.B., Foster, G.L., Erez, J., Prentice, K.C., Kucera, M., Bostock,
 764 H.C., Martínez-Botí, M.A., Milton, J.A., Wilson, P.A., Marshall, B.J., Elliott, T.,
 765 2013. Calibration of the boron isotope proxy in the planktonic foraminifera
 766 *Globigerinoides ruber* for use in palaeo-CO₂ reconstruction. *Earth and Planetary*
 767 *Science Letters*, 364: 111-122.

768 Holcomb, M., Cohen, A.L., Gabitov, R.I., Hutter, J.L., 2009. Compositional and
 769 morphological features of aragonite precipitated experimentally from seawater
 770 and biogenically by corals. *Geochimica et Cosmochimica Acta*, 73: 4166-4179.

771 Holcomb, M., Cohen, A.L., McCorkle, D.C., 2013. An evaluation of staining techniques
 772 for marking daily growth in scleractinian corals. *Journal of Experimental Marine*
 773 *Biology and Ecology*, 440: 126-131.

774 Holcomb, M., Rankenburg, K., McCulloch, M., 2014. High Precision MC-ICP-MS
 775 measurements of $\delta^{11}\text{B}$: matrix effects in direct injection and spray chamber
 776 sample introduction systems, in Grice, K., (Ed.) *Principles and Practice of*
 777 *Analytical Techniques in Geosciences*. Royal Society of Chemistry, England:
 778 251-270.

779 Hong, S.C., Kim, M.-S., Chung, J.G., 2002. Adsorption characteristics of Pb(II) on
 780 calcite-type calcium carbonate by batch and continuous reactors. *Journal of*
 781 *Industrial Engineering and Chemistry*, 8: 305-312.

782 Ihlenfeld, C., Norman, M.D., Gagan, M.K., Drysdale, R.N., Maas, R., Webb, J., 2003.
 783 Climatic significance of seasonal trace element and stable isotope variations in a
 784 modern freshwater tufa. *Geochimica et Cosmochimica Acta*, 67(13): 2341-2357.

785 Ingalls, A.E., Lee, C., Druffel, E.R.M., 2003. Preservation of organic matter in mound-
 786 forming coral skeletons. *Geochimica et Cosmochimica Acta*, 67: 2827-2841.

787 Inoue, M., Nohara, M., Okai, T., Suzuki, A., Kawahata, H., 2004. Concentrations of
 788 Trace Elements in Carbonate Reference Materials Coral JCp-1 and Giant Clam
 789 JCt-1 by Inductively Coupled Plasma-Mass Spectrometry. *Geostandards and*
 790 *Geoanalytical Research*, 28(3): 411-416.

791 Inoue, M., Suzuki, A., Nohara, M., Kawahata, H., 2009. Assessment of marine pollution
 792 in tropical and subtropical region using trace elements in coral skeletons. In:
 793 Davin, T.B., Brannet, A.P. (Eds.), *Coral Reefs: Biology, Threats and Restoration*.
 794 Nova Science Publishers, Inc., New York, pp. 183-208.

795 Kitano, Y., Okumura, M., Idogaki, M., 1978. Coprecipitation of borate-boron with
 796 calcium carbonate. *Geochemical Journal*, 12: 183-189.

797 Klinkhammer, G.P., Haley, B.A., Mix, A.C., Benway, H.M., Cheseby, M., 2004.
 798 Evaluation of automated flow-through time-resolved analysis of foraminifera for
 799 Mg/Ca paleothermometry. *Paleoceanography*, 19(4): PA4030.

800 Klochko, K., Kaufman, A.J., Yao, W., Byrne, R.H., Tossell, J.A., 2006. Experimental
801 measurement of boron isotope fractionation in seawater. *Earth and Planetary*
802 *Science Letters*, 248(1-2): 276-285.

803 Krause-Nehring, J., Klügel, A., Nehrke, G., Brellochs, B., Brey, T., 2011. Impact of
804 sample pretreatment on the measured element concentrations in the bivalve
805 *Arctica islandica*. *Geochemistry, Geophysics, Geosystems*, 12(7): Q07015,
806 doi:10.1029/2011GC003630.

807 LaVigne, M., Field, M.P., Anagnostou, E., Grottoli, A.G., Wellington, G.M., Sherrell,
808 R.M., 2008. Skeletal P/Ca tracks upwelling in Gulf of Panamá coral: Evidence for
809 a new seawater phosphate proxy. *Geophysical Research Letters*, 35(5): L05604.

810 Lea, D.W., Boyle, E.A., 1993. Determinations of carbonate-bound barium in foraminifera
811 and corals by isotope dilution plasma-mass spectrometry. *Chemical Geology*,
812 103: 73-84.

813 Leder, J.J., Swart, P.K., Szmant, A.M., Dodge, R.E., 1996. The origin of variations in the
814 isotopic record of scleractinian corals: I. Oxygen. *Geochimica et Cosmochimica*
815 *Acta*, 60: 2857-2870.

816 Macintyre, I.G., Towe, K.M., Houck, J.E., Buddemeier, R.W., Chave, K.E., 1976.
817 Skeletal calcite in living scleractinian corals: Microboring fillings, not primary
818 skeletal deposits. *Science*, 193(4254): 701-702.

819 McConchie, D., Harriott, V.J., 1992. The partitioning of metals between tissue and
820 skeletal parts of corals: application in pollution monitoring. *Proceedings of the 7th*
821 *International Coral Reef Symposium*, 1: 97-103.

822 McCulloch, M., Falter, J., Trotter, J., Montagna, P., 2012. Coral resilience to ocean
823 acidification and global warming through pH up-regulation. *Nature Climate*
824 *Change*, 2(8): 623-627.

825 McCulloch, M.T., Holcomb, M., Rankenburg, K., Trotter, J.A., 2014. Rapid, high-
826 precision measurements of boron isotopic compositions in marine carbonates.
827 *Rapid Communications in Mass Spectrometry*, 28(24): 2704-12.

828 Millero, F., Huang, F., Zhu, X., Liu, X., Zhang, J.-Z., 2001. Adsorption and desorption of
829 phosphate on calcite and aragonite in seawater. *Aquatic Geochemistry*, 7(1): 33-
830 56.

831 Millero, F.J., Woosley, R., DiTrollo, B., Waters, J., 2009. Effect of ocean acidification on
832 the speciation of metals in seawater. *Oceanography*, 22: 72-85.

833 Mitsuguchi, T., Kawakami, T., 2012. Potassium and other minor elements in *Porites*
834 corals: implications for skeletal geochemistry and paleoenvironmental
835 reconstruction. *Coral Reefs*, 31(3): 671-681.

836 Mitsuguchi, T., Uchida, T., Matsumoto, E., Isdale, P.J., Kawana, T., 2001. Variations in
837 Mg/Ca, Na/Ca, and Sr/Ca ratios of coral skeletons with chemical treatments:
838 implications for carbonate geochemistry. *Geochimica et Cosmochimica Acta*,
839 65(17): 2865-2874.

840 Mohnen, D., 2008. Pectin structure and biosynthesis. *Current Opinion in Plant Biology*,
841 11(3): 266-77.

842 Montagna, P., McCulloch, M., Mazzoli, C., Silenzi, S., Odorico, R., 2007. The non-
843 tropical coral *Cladocora caespitosa* as the new climate archive for the

844 Mediterranean: high-resolution (~weekly) trace element systematics. *Quaternary*
845 *Science Reviews*, 26(3-4): 441-462.

846 Nagtegaal, R., Grove, C.A., Kasper, S., Zinke, J., Boer, W., Brummer, G.-J.A., 2012.
847 Spectral luminescence and geochemistry of coral aragonite: Effects of whole-core
848 treatment. *Chemical Geology*, 318-319: 6-15.

849 Nothdurft, L.D., Webb, G.E., Buster, N.A., Holmes, C.W., Sorauf, J.E., Kloprogge, J.T.,
850 2005. Brucite microbialites in living coral skeletons: Indicators of extreme
851 microenvironments in shallow-marine settings. *Geology*, 33(3): 169-172.

852 Okai, T., Suzuki, A., Kawahata, H., Terashima, S., Imai, N., 2002. Preparation of a new
853 Geological Survey of Japan Geochemical Reference Material: Coral JCp-1.
854 *Geostandards Newsletter*, 26(1): 95-99.

855 Pak, D.K., Lea, D.W., Kennett, J.P., 2004. Seasonal and interannual variation in Santa
856 Barbara Basin water temperatures observed in sediment trap foraminiferal Mg/Ca.
857 *Geochemistry, Geophysics, Geosystems*, 5(12): Q12008.

858 Pingitore, N.E., Fretzdorff, S.B., Seitz, B.P., Estrada, L.Y., Borrego, P.M., Crawford,
859 G.M., Love, K.M., 1993. Dissolution kinetics of CaCO₃ in common laboratory
860 solvents. *Journal of Sedimentary Research*, 63(4): 641-645.

861 Reichelt-Brushett, A.J., McOrist, G., 2003. Trace metals in the living and nonliving
862 components of scleractinian corals. *Marine Pollution Bulletin*, 46(12): 1573-1582.

863 Reynaud, S., Ferrier-Pages, C., Meibom, A., Mostefaoui, S., Mortlock, R., Fairbanks, R.,
864 Allemand, D., 2007. Light and temperature effects on Sr/Ca and Mg/Ca ratios in
865 the scleractinian coral *Acropora sp.* *Geochimica et Cosmochimica Acta*, 71: 354-
866 362.

867 Robinson, L.F., Belshaw, N.S., Henderson, G.M., 2004. U and Th concentrations and
 868 isotope ratios in modern carbonates and waters from the Bahamas. *Geochimica et*
 869 *Cosmochimica Acta*, 68(8): 1777-1789.

870 Rosenthal, Y., Boyle, E.A., Labeyrie, L., 1997. Last Glacial Maximum paleochemistry
 871 and deepwater circulation in the Southern Ocean: Evidence from foraminiferal
 872 cadmium. *Paleoceanography*, 12(6): 787-796.

873 Rouff, A.A., Elzinga, E.J., Reeder, R.J., Fisher, N.S., 2005. The influence of pH on the
 874 kinetics, reversibility and mechanisms of Pb(II) sorption at the calcite-water
 875 interface. *Geochimica et Cosmochimica Acta*, 69(22): 5173-5186.

876 Saenger, C., Cohen, A.L., Oppo, D.W., Hubbard, D., 2008. Interpreting sea surface
 877 temperature from strontium/calcium ratios in *Montastrea* corals: Link with growth
 878 rate and implications for proxy reconstructions. *Paleoceanography*, 23(3):
 879 PA3102.

880 Sanyal, A., Hemming, N.G., Broecker, W.S., Lea, D.W., Spero, H.J., Hanson, G.N.,
 881 1996. Oceanic pH control on the boron isotopic composition of foraminifera:
 882 Evidence from culture experiments. *Paleoceanography*, 11(5): 513-517.

883 Semião, A.J.C., Rossiter, H.M.A., Schäfer, A.I., 2010. Impact of organic matter and
 884 speciation on the behaviour of uranium in submerged ultrafiltration. *Journal of*
 885 *Membrane Science*, 348(1-2): 174-180.

886 Shen, G.T., Boyle, E.A., 1988. Determination of lead, cadmium and other trace metals in
 887 annually-banded corals. *Chemical Geology*, 67(1-2): 47-62.

888 Shirai, K., Sowa, K., Watanabe, T., Sano, Y., Nakamura, T., Clode, P., 2012.
889 Visualization of sub-daily skeletal growth patterns in massive *Porites* corals
890 grown in Sr-enriched seawater. *Journal of Structural Biology*, 180(1): 47-56.

891 Sinclair, D.J., 2005. Non-river flood barium signals in the skeletons of corals from
892 coastal Queensland, Australia. *Earth and Planetary Science Letters*, 237(3-4):
893 354-369.

894 Sinclair, D.J., Kinsley, L.P.J., McCulloch, M.T., 1998. High resolution analysis of trace
895 elements in corals by laser ablation ICP-MS. *Geochimica et Cosmochimica Acta*,
896 62: 1889-1901.

897 Spivack, A.J., Edmond, J.M., 1986. Determination of boron isotope ratios by thermal
898 ionization mass spectrometry of the dicesium metaborate cation. *Analytical*
899 *Chemistry*, 58(1): 31-35.

900 St. John, B.E., 1974. Heavy metals in the skeletal carbonate of scleractinian corals.
901 *Proceedings of the 2nd International Coral Reef Symposium*, 2: 461-469.

902 Stoll, H.M., Encinar, J.R., Alonso, J.I.G., 2001. A first look at paleotemperature
903 prospects from Mg in coccolith carbonate: cleaning techniques and culture
904 measurements. *Geochemistry, Geophysics, Geosystems*, 2: 144.

905 Tambutté, E., Allemand, D., Zoccola, D., Meibom, A., Lotto, S., Caminiti, N., Tambutté,
906 S., 2007. Observations of the tissue-skeleton interface in the scleractinian coral
907 *Stylophora pistillata*. *Coral Reefs*, 26(3): 517-529.

908 Tatsumoto, M., Goldberg, E.D., 1959. Some aspects of the marine geochemistry of
909 uranium. *Geochimica et Cosmochimica Acta*, 17: 201-208.

910 Tessier, A., Campbell, P.G.C., Bisson, M., 1979. Sequential extraction procedure for the
 911 speciation of particulate trace metals. *Analytical Chemistry*, 51(7): 844-851.
 912 Tudhope, A.W., Lea, D.W., Shimmield, G.B., Chilcott, C.P., Head, S., 1996. Monsoon
 913 climate and Arabian Sea coastal upwelling recorded in massive corals from
 914 southern Oman. *Palaios*, 11: 347-361.
 915 Turner, D.R., Whitfield, M., Dickson, A.G., 1981. The equilibrium speciation of
 916 dissolved components in freshwater and sea water at 25°C and 1 atm pressure.
 917 *Geochimica et Cosmochimica Acta*, 45(6): 855-881.
 918 van de Flierdt, T., Robinson, L.F., Adkins, J.F., 2010. Deep-sea coral aragonite as a
 919 recorder for the neodymium isotopic composition of seawater. *Geochimica et*
 920 *Cosmochimica Acta*, 74(21): 6014-6032.
 921 Vengosh, A., Kolodny, Y., Starinsky, A., Chivas, A.R., McCulloch, M.T., 1991.
 922 Coprecipitation and isotopic fractionation of boron in modern biogenic
 923 carbonates. *Geochimica et Cosmochimica Acta*, 55: 2901-2910.
 924 Wang, Z., Gaetani, G., Liu, C., Cohen, A., 2013a. Oxygen isotope fractionation between
 925 aragonite and seawater: Developing a novel kinetic oxygen isotope fractionation
 926 model. *Geochimica et Cosmochimica Acta*, 117: 232-251.
 927 Wang, Z., Hu, P., Gaetani, G., Liu, C., Saenger, C., Cohen, A., Hart, S., 2013b.
 928 Experimental calibration of Mg isotope fractionation between aragonite and
 929 seawater. *Geochimica et Cosmochimica Acta*, 102: 113-123.
 930 Wara, M.W., Delaney, M.L., Bullen, T.D., Ravelo, A.C., 2003. Possible roles of pH,
 931 temperature, and partial dissolution in determining boron concentration and
 932 isotopic composition in planktonic foraminifera. *Paleoceanography*, 18(4): 1100.

933 Watanabe, T., Minagawa, M., Oba, T., Winter, A., 2001. Pretreatment of coral aragonite
 934 from Mg and Sr analysis: implications for coral thermometers. *Geochemical*
 935 *Journal*, 35: 265-269.

936 Wierzbowski, H., 2007. Effects of pre-treatments and organic matter on oxygen and
 937 carbon isotope analyses of skeletal and inorganic calcium carbonate. *International*
 938 *Journal of Mass Spectrometry*, 268(1): 16-29.

939 Yoshioka, S., Ohde, S., Kitano, Y., Kanamori, N., 1986. Behaviour of magnesium and
 940 strontium during the transformation of coral aragonite to calcite in aquatic
 941 environments. *Marine Chemistry*, 18(1): 35-48.

942 Young, S.D., 1971. Organic material from scleractinian coral skeletons - I. Variation in
 943 composition between several species. *Comparative Biochemistry and*
 944 *Physiology*, 40B: 113-120.

945 Zinke, J., Rountrey, A., Feng, M., Xie, S.P., Dissard, D., Rankenburg, K., Lough, J.M.,
 946 McCulloch, M.T., 2014. Corals record long-term Leeuwin current variability
 947 including Ningaloo Niño/Niña since 1795. *Nature Communications*, 5.
 948
 949
 950

Supplemental materials

Synthetic aragonite

Alkalinity and pH samples were taken periodically over the course of the aragonite growth experiments. Alkalinity samples were diluted immediately with a ~0.68 M NaCl solution. Measurements were made as described by Holcomb et al. (2012). Alkalinity, pH, predicted quantities of aragonite precipitated, and the times of isotope spikes are plotted in Figure S1. A scanning electron microscopy (SEM) image of the resultant precipitate is shown in Figure S6.

Supplemental results and discussion

In treatments of the TS3 aragonite powder conducted at UWA, no significant differences were found among rinsing, bleaching, and NaOH for any of the measured ratios other than U/Ca ($F_{2,6} = 14.4$, $p < 0.01$). Bleached samples had higher U/Ca than rinsed ($p < 0.01$) and NaOH ($p < 0.05$) treated samples (Fig. S4). For the same sample treated at WHOI, there was no significant effect of NaOH concentration between 0.006 M and 0.1 M on Mg/Ca, Sr/Ca, Ba/Ca, or U/Ca. However, bleach significantly increased Sr/Ca ($F_{4,17} = 43.1$, bleach higher than other treatments $p < 0.001$) and decreased U/Ca ($F_{4,17} = 32.3$, bleach lower than other treatments $p < 0.001$) relative to cleaning with H₂O or any of the NaOH treatments. (Figure S5).

Although both bleach (Figs. 2,3,4,S5) and NaOH (see results, section 3.2.2) were associated with declines in the U/Ca ratio, this was not always observed (Fig. S4), even in treatments of the same sample (Figs. S4,S5). This suggests that the reduction in U/Ca is not related to the high pH of the wash solution alone, but due to some other factor. Variation in the CO_3^{2-} content of the solution (due to variable uptake of CO_2) is suggested as a potential explanation (see discussion section 4.2). In one instance, treatment with bleach was associated with an unexpected increase in Sr/Ca (Fig. S5). Although it is not known why bleach might result in an increase in Sr/Ca, it could be related to Sr present in the bleach solution used. Krause-Nehring et al. (2011) measured the composition of the bleach solution used in their experiments and found high Sr/Ca ratios (as well as contamination with other elements), suggesting that bleach has the potential to contaminate samples. Nagtegaal et al., (2012) also reported differing effects of bleach obtained from different suppliers. Hydrogen peroxide too has the potential to contaminate samples. For a synthetic aragonite sample (1/19/11 d), treatment with H_2O_2 was associated with an increase in Sn/Ca, which persisted despite repeated rinsing with H_2O post treatment (data not shown). Thus care must be taken in selecting reagents for cleaning samples to avoid introducing new sources of contamination.

Supplemental References

Arslan, Z., Secor, D.H., 2008. High resolution micromill sampling for analysis of fish otoliths by ICP-MS: effects of sampling and specimen preparation on trace element fingerprints. *Marine Environmental Research*, 66(3): 364-71.

996 Bian, N., Martin, P.A., 2010. Investigating the fidelity of Mg/Ca and other elemental data
 997 from reductively cleaned planktonic foraminifera. *Paleoceanography*, 25(2),
 998 PA2215, doi:10.1029/2009PA001796.

999 Boiteau, R., Greaves, M., Elderfield, H., 2012. Authigenic uranium in foraminiferal
 1000 coatings: A proxy for ocean redox chemistry. *Paleoceanography*, 27(3): PA3227.

1001 Campana, S.E., Chouinard, G.A., Hanson, J.M., Fréchet, A., Bratney, J., 2000. Otolith
 1002 elemental fingerprints as biological tracers of fish stocks. *Fisheries Research*,
 1003 46(1–3): 343-357.

1004 Davies, C.A., Brophy, D., Jeffries, T., Gosling, E., 2011. Trace elements in the otoliths
 1005 and dorsal spines of albacore tuna (*Thunnus alalunga*, Bonnaterre, 1788): An
 1006 assessment of the effectiveness of cleaning procedures at removing postmortem
 1007 contamination. *Journal of Experimental Marine Biology and Ecology*, 396(2):
 1008 162-170.

1009 Delaney, M.L., Linn, L.J., Davies, P.J., 1996. Trace and minor element ratios in
 1010 Halimeda aragonite from the Great Barrier Reef. *Coral Reefs*, 15(3): 181-189.

1011 Duplessy, J.C., Lalou, C., Vinot, A.C., 1970. Differential isotopic fractionation in benthic
 1012 foraminifera and paleotemperatures reassessed. *Science*, 168(3928): 250-251.

1013 Durazzi, J.T., 1977. Stable isotopes in the ostracod shell: a preliminary study.
 1014 *Geochimica et Cosmochimica Acta*, 41(8): 1168-1170.

1015 Emiliani, C., 1966. Paleotemperature analysis of Caribbean cores P6304-8 and P6304-9
 1016 and a generalized temperature curve for the past 425,000 years. *The Journal of*
 1017 *Geology*, 74: 109-126.

1018 Erez, J., Honjo, S., 1981. Comparison of isotopic composition of planktonic foraminifera
 1019 in plankton tows, sediment traps and sediments. *Palaeogeography*,
 1020 *Palaeoclimatology, Palaeoecology*, 33: 129-156.

1021 Fallet, U., Boer, W., van Assen, C., Greaves, M., Brummer, G.-J.A., 2009. A novel
 1022 application of wet oxidation to retrieve carbonates from large organic-rich
 1023 samples for ocean-climate research. *Geochemistry, Geophysics, Geosystems*,
 1024 10(8): Q08004, doi:10.1029/2009GC002573.

1025 Feldmeijer, W., Metcalfe, B., Scussolini, P., Arthur, K., 2013. The effect of chemical
 1026 pretreatment of sediment upon foraminiferal-based proxies. *Geochemistry*,
 1027 *Geophysics, Geosystems*, 14(10): 3996-4014.

1028 Gaffey, S.J., Kolak, J.J., Bronnimann, C.E., 1991. Effects of drying, heating, annealing,
 1029 and roasting on carbonate skeletal material, with geochemical and diagenetic
 1030 implications. *Geochimica et Cosmochimica Acta*, 55: 1627-1640.

1031 Ganssen, G., 1981. Isotopic analysis of foraminifera shells: Interference from chemical
 1032 treatment. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 33(1–3): 271-
 1033 276.

1034 Glock, N., Eisenhauer, A., Liebetrau, V., Wiedenbeck, M., Hensen, C., Nehrke, G., 2012.
 1035 EMP and SIMS studies on Mn/Ca and Fe/Ca systematics in benthic foraminifera
 1036 from the Peruvian OMZ: a contribution to the identification of potential redox
 1037 proxies and the impact of cleaning protocols. *Biogeosciences*, 9(1): 341-359.

1038 Gray, W., Holmes, J., Shevenell, A., 2014. Evaluation of foraminiferal trace element
 1039 cleaning protocols on the Mg/Ca of marine ostracod genus *Krithe*. *Chemical*
 1040 *Geology*, 382(0): 14-23.

1041 Grossman, E.L., Betzer, P.R., Dudley, W.C., Dunbar, R.B., 1986. Stable isotopic
1042 variation in pteropods and atlantids from North Pacific sediment traps. *Marine*
1043 *Micropaleontology*, 10: 9-22.

1044 Hastings, D.W., Russell, A.D., Emerson, S.R., 1998. Foraminiferal magnesium in
1045 *Globeriginoidea sacculifer* as a paleotemperature proxy. *Paleoceanography*,
1046 13(2): 161-169.

1047 Holcomb, M., Cohen, A.L., McCorkle, D.C., 2012. An investigation of the calcification
1048 response of the scleractinian coral *Astrangia poculata* to elevated pCO₂ and the
1049 effects of nutrients, zooxanthellae and gender. *Biogeosciences*, 9(1): 29-39.

1050 Holmes, J.A., 1992. Trace-element chemistry of non-marine ostracod shells: a
1051 preliminary evaluation of cleaning methods. *Journal of Micropalaeontology*,
1052 11(1): 36.

1053 Inoue, M., Suzuki, A., Nohara, M., Kan, H., Edward, A., Kawahata, H., 2004a. Coral
1054 skeletal tin and copper concentrations at Pohnpei, Micronesia: possible index for
1055 marine pollution by toxic anti-biofouling paints. *Environmental Pollution*, 129(3):
1056 399-407.

1057 Jin, Z., Bickle, M., Chapman, H., Yu, J., Greaves, M., Wang, S., Chen, S., 2006. An
1058 experimental evaluation of cleaning methods for fossil ostracod Mg/Ca and Sr/Ca
1059 determination. *Journal of Paleolimnology*, 36(2): 211-218.

1060 Keatings, K.W., Holmes, J.A., Heaton, T.H.E., 2006. Effects of pre-treatment on ostracod
1061 valve chemistry. *Chemical Geology*, 235: 250-261.

1062 Kontrovitz, M., Slack, J.M., Yuhong, Z., 1991. On the use of some phosphates in the
1063 preparation of ostracod shells. *Journal of Micropalaeontology*, 10(2): 121-126.

1064 Land, L.S., Lang, J.C., Barnes, D.J., 1975. Extension rate: a primary control on the
 1065 isotopic composition of West Indian (Jamaican) scleractinian reef coral skeletons.
 1066 Marine Biology, 33: 221-233.

1067 LaVigne, M., Hill, T.M., Sanford, E., Gaylord, B., Russell, A.D., Lenz, E.A., Hosfelt,
 1068 J.D., Young, M.K., 2013. The elemental composition of purple sea urchin
 1069 (*Strongylocentrotus purpuratus*) calcite and potential effects of pCO₂ during early
 1070 life stages. Biogeosciences, 10(6): 3465-3477.

1071 Lécuyer, C., 1996. Effects of heating on the geochemistry of biogenic carbonates.
 1072 Chemical Geology, 129(3–4): 173-183.

1073 Li, X., Weiguo, L., Pengxi, Z., Zhisheng, A., Ling, Z., 2007. Species, valve size, and
 1074 pretreatment effects on $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of ostracod valves from Lake
 1075 Qinghai, Qinghai–Tibet Plateau. Chemical Geology, 246(1–2): 124-134.

1076 Linn, L.J., Delaney, M.L., Druffel, E.R.M., 1990. Trace metals in contemporary and
 1077 seventeenth-century Galapagos coral: Records of seasonal and annual variations.
 1078 Geochimica et Cosmochimica Acta, 54(2): 387-394.

1079 Love, K.M., Woronow, A., 1991. Chemical changes induced in aragonite using
 1080 treatments for the destruction of organic material. Chemical Geology, 93(3–4):
 1081 291-301.

1082 Marr, J.P., Bostock, H.C., Carter, L., Bolton, A., Smith, E., 2013. Differential effects of
 1083 cleaning procedures on the trace element chemistry of planktonic foraminifera.
 1084 Chemical Geology, 351: 310-323.

1085 Martin, P.A., Lea, D.W., 2002. A simple evaluation of cleaning procedures on fossil
 1086 benthic foraminiferal Mg/Ca. *Geochemistry, Geophysics, Geosystems*, 3(10):
 1087 8401.

1088 McConnaughey, T., 1989. ^{13}C and ^{18}O isotopic disequilibrium in biological carbonates: I.
 1089 Patterns. *Geochimica et Cosmochimica Acta*, 53: 151-162.

1090 Mischke, S., Zhang, C., Börner, A., 2008. Bias of ostracod stable isotope data caused by
 1091 drying of sieve residues from water. *Journal of Paleolimnology*, 40(1): 567-575.

1092 Misra, S., Greaves, M., Owen, R., Kerr, J., Elmore, A.C., Elderfield, H., 2014a.
 1093 Determination of B/Ca of natural carbonates by HR-ICP-MS. *Geochemistry,*
 1094 *Geophysics, Geosystems*, 15(4): 1617-1628.

1095 Misra, S., Owen, R., Kerr, J., Greaves, M., Elderfield, H., 2014b. Determination of $\delta^{11}\text{B}$
 1096 by HR-ICP-MS from mass limited samples: Application to natural carbonates and
 1097 water samples. *Geochimica et Cosmochimica Acta*, 140: 531-552.

1098 Mook, W.G., 1971. Paleotemperatures and chlorinities from stable carbon and oxygen
 1099 isotopes in shell carbonate. *Palaeogeography, Palaeoclimatology, Palaeoecology*,
 1100 9: 245-263.

1101 Pena, L.D., Calvo, E., Cacho, I., Eggins, S., Pelejero, C., 2005. Identification and
 1102 removal of Mn-Mg-rich contaminant phases on foraminiferal tests: Implications
 1103 for Mg/Ca past temperature reconstructions. *Geochemistry, Geophysics,*
 1104 *Geosystems*, 6(9): Q09P02.

1105 Penman, D.E., Hönisch, B., Rasbury, E.T., Hemming, N.G., Spero, H.J., 2013. Boron,
 1106 carbon, and oxygen isotopic composition of brachiopod shells: Intra-shell

1107 variability, controls, and potential as a paleo-pH recorder. *Chemical Geology*,
 1108 340: 32-39.

1109 Roberts, N.L., Piotrowski, A.M., Elderfield, H., Eglinton, T.I., Lomas, M.W., 2012. Rare
 1110 earth element association with foraminifera. *Geochimica et Cosmochimica Acta*,
 1111 94(0): 57-71.

1112 Saenger, C., Affek, H.P., Felis, T., Thiagarajan, N., Lough, J.M., Holcomb, M., 2012.
 1113 Carbonate clumped isotope variability in shallow water corals: Temperature
 1114 dependence and growth-related vital effects. *Geochimica et Cosmochimica Acta*,
 1115 99: 224-242.

1116 Saenger, C., Wang, Z., Gaetani, G., Cohen, A., Lough, J.M., 2013. The influence of
 1117 temperature and vital effects on magnesium isotope variability in *Porites* and
 1118 *Astrangia* corals. *Chemical Geology*, 360–361(0): 105-117.

1119 Sarkar, A., Ramesh, R., Bhattacharya, S.K., 1990. Effect of sample pretreatment and size
 1120 fraction on the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of foraminifera in Arabian Sea sediments.
 1121 *Terra Nova*, 2(5): 488-493.

1122 Serrano, O., Serrano, L., Mateo, M.A., 2008. Effects of sample pre-treatment on the $\delta^{13}\text{C}$
 1123 and $\delta^{18}\text{O}$ values of living benthic foraminifera. *Chemical Geology*, 257(3-4): 218-
 1124 220.

1125 Shen, G., Campbell, T., Dunbar, R., Wellington, G., Colgan, M., Glynn, P., 1991.
 1126 Paleochemistry of manganese in corals from the Galapagos Islands. *Coral Reefs*,
 1127 10(2): 91-100.

1128 Shen, G.T., Dunbar, R.B., 1995. Environmental controls on uranium in reef corals.
 1129 *Geochimica et Cosmochimica Acta*, 59(10): 2009-2024.

1130 Uchikawa, J., Penman, D.E., Zachos, J.C., Zeebe, R.E., 2015. Experimental evidence for
 1131 kinetic effects on B/Ca in synthetic calcite: Implications for potential B(OH)_4^-
 1132 and B(OH)_3 incorporation. *Geochimica et Cosmochimica Acta*, 150: 171-191.
 1133 Vetter, L., Spero, H.J., Russell, A.D., Fehrenbacher, J.S., 2013. LA-ICP-MS depth
 1134 profiling perspective on cleaning protocols for elemental analyses in planktic
 1135 foraminifers. *Geochemistry, Geophysics, Geosystems*, 14(8): 2916-2931.
 1136 Waelbroeck, C., Mulitza, S., Spero, H., Dokken, T., Kiefer, T., Cortijo, E., 2005. A
 1137 global compilation of late Holocene planktonic foraminiferal $\delta^{18}\text{O}$: relationship
 1138 between surface water temperature and $\delta^{18}\text{O}$. *Quaternary Science Reviews*, 24(7-
 1139 9): 853-868.
 1140 Wang, X.T., Sigman, D.M., Cohen, A.L., Sinclair, D.J., Sherrell, R.M., Weigand, M.A.,
 1141 Erler, D.V., Ren, H., 2015. Isotopic composition of skeleton-bound organic
 1142 nitrogen in reef-building symbiotic corals: A new method and proxy evaluation at
 1143 Bermuda. *Geochimica et Cosmochimica Acta*, 148: 179-190.
 1144 Weldeab, S., Schneider, R.R., Kölling, M., 2006. Comparison of foraminiferal cleaning
 1145 procedures for Mg/Ca paleothermometry on core material deposited under
 1146 varying terrigenous-input and bottom water conditions. *Geochemistry,*
 1147 *Geophysics, Geosystems*, 7(4), Q04P12, doi:10.1029/2005GC000990.
 1148 Wotte, T., Shields-Zhou, G.A., Strauss, H., 2012. Carbonate-associated sulfate:
 1149 Experimental comparisons of common extraction methods and recommendations
 1150 toward a standard analytical protocol. *Chemical Geology*, 326-327: 132-144.

1151 Yu, J., Elderfield, H., Greaves, M., Day, J., 2007. Preferential dissolution of benthic
1152 foraminiferal calcite during laboratory reductive cleaning. *Geochemistry,*
1153 *Geophysics, Geosystems*, 8(6), Q06016, doi:10.1029/2006GC001571.
1154

Supplemental tables and figures

Table S1. Studies examining the effects of different cleaning procedures on calcium carbonate samples. Independent treatments that were compared are separated by “,”, mixed reagents are indicated by “/” between reagent names. For studies which compared multi-step procedures instead of individual treatments, major steps are listed with “+” between each step. Only the major treatment compared is listed in the cleaning agents column, some studies used an initial pre-treatment prior to comparing specific reagents. Studies are ordered from most recent to earliest.

Cleaning agents	Sample type	Measurements	Study
rinsing, H ₂ O ₂ , HNO ₃ , NaOH, NaOCl	abiotic and coral aragonite, foraminifera calcite	$\delta^{11}\text{B}$, Li, B, Na, Mg, Sr, Ba, Pb, U	current study
H ₂ O ₂ /NaOH + HNO ₃ + hydrazine/ammonia/citrate	abiotic calcite	B	Uchikawa et al., 2015
NaOCl, persulfate	coral aragonite	$\delta^{15}\text{N}$	Wang et al., 2015
rinsing and methanol, H ₂ O ₂ /NaOH, hydrazine/ammonia/citrate	ostracod calcite	Mg, Al, Mn, Fe, Sr, microscopy	Gray et al., 2014
H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/acetate, combinations thereof	foraminifera calcite	B	Misra et al., 2014a
H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/acetate, combinations thereof	foraminifera calcite	$\delta^{11}\text{B}$	Misra et al., 2014b
H ₂ O ₂ , H ₂ O ₂ /(NaPO ₃) ₆ , Na ₄ P ₂ O ₇	foraminifera calcite sediments	pH, fragmentation, $\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Mg	Feldmeijer et al., 2013
rinsing, H ₂ O ₂ /NaOH, NaOCl, heat, combinations thereof	sea urchin calcite	Mg, Sr	LaVigne et al., 2013
H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/citrate	foraminifera calcite	Mg, Al, Mn, Zn, Sr, Ba	Marr et al., 2013
NaOCl, heat + NaOCl	brachiopod	$\delta^{11}\text{B}$	Penman et al.,

	calcite		2013
methanol, acetate	coral aragonite	$\delta^{25}\text{Mg}$, $\delta^{26}\text{Mg}$	Saenger et al., 2013
$\text{H}_2\text{O}_2/\text{NaOH}$ + hydrazine/ammonia/citrate + DTPA/NaOH	foraminifera calcite	Mg, Ba	Vetter et al., 2013
rinsing, methanol, acetate	abiotic aragonite	$\delta^{26}\text{Mg}$, Mg, Sr, Ba, U	Wang et al., 2013b
rinsing, $\text{H}_2\text{O}_2/\text{NaOH}$, NaOCl, $\text{NH}_2\text{OH}\cdot\text{HCl}/\text{NH}_4\text{OH}$	abiotic and coccolith calcite, organics	Mg, P, Ca, Fe, Sr	Blanco-Ameijeiras et al., 2012
$\text{H}_2\text{O}_2/\text{NaOH}$, HNO_3 , hydrazine/ammonia/citrate, combinations thereof	foraminifera calcite	U	Boiteau et al., 2012
H_2O + methanol + $\text{H}_2\text{O}_2/\text{NaOH}$ + HNO_3	foraminifera calcite	Mg, S, Ca, Mn, Fe	Glock et al., 2012
rinsing, H_2O_2 , HNO_3	coral aragonite	Na, Mg, K, Sr	Mitsuguchi and Kawakami, 2012
H_2O_2 , HNO_3 , NaOCl	coral aragonite (slabs)	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Mg, Sr, Y, Ba, U, luminescence	Nagtegaal et al., 2012
rinsing, $\text{H}_2\text{O}_2/\text{NaOH}$, hydrazine/ammonia/citrate, DTPA/NaOH	foraminifera calcite	Nd	Roberts et al., 2012
H_2O_2	coral aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Δ_{47}	Saenger et al., 2012
rinsing, H_2O_2 , NaOCl, NaCl	fossil calcium carbonates	$\delta^{34}\text{S}$, S	Wotte et al., 2012
rinsing, H_2O_2 , HNO_3	otolith aragonite	Mg, Mn, Sr, Ba	Davies et al., 2011
rinsing, H_2O_2 , NaOH, NaOCl, acetone	bivalve aragonite, abiotic calcite	N, Mg, Ca, Mn, Sr, Ba, crystallography	Krause-Nehring et al., 2011
citrate, hydrazine, NH_4OH , and mixtures thereof	foraminifera calcite	Na, Mg, Al, Mn, Fe, Sr, Cd, Ba, La, Ce, Nd, U	Bian and Martin, 2010
physical cleaning, $\text{H}_2\text{O}_2/\text{NaOH}$ + $\text{H}_2\text{O}_2/\text{HClO}_4$, methanol + HNO_3 , hydrazine/ammonia/citrate, EDTA + HNO_3	coral aragonite	Nd isotopes, Nd, Th, U	van de Flierdt et al., 2010
rinsing, H_2O_2 ,	foraminifera	pH, $\delta^{13}\text{C}$, $\delta^{18}\text{O}$,	Fallet et al., 2009

H ₂ O ₂ /NaOH, heat	calcite and bivalve aragonite sediments	Mg, Fe, Sr, Ba	
rinsing, H ₂ O ₂ /NaOH, HNO ₃	coral aragonite	Cu, Sn	Inoue et al., 2009
rinsing/H ₂ O ₂ /HNO ₃	otolith aragonite	Al, V, Mn, Co, Ni, Cu, Zn, Ga, Cd, Pb, Bi	Arslan and Secor 2008
rinsing, ethanol (effect of evaporation)	ostracod calcite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$, microscopy	Mischke et al., 2008
H ₂ O ₂ , NaOCl, heat, ethanol, methanol, sonication, + combinations thereof	foraminifera calcite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Serrano et al., 2008
rinsing + methanol, H ₂ O ₂ , methanol	ostracod calcite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Li et al., 2007
H ₂ O ₂ , NaOCl, heat	abiotic and biotic calcite and aragonite, organics	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Wierzbowski, 2007
rinsing, H ₂ O ₂ /NaOH, NaOH, NH ₃ , hydrazine, citrate, and mixtures thereof	foraminifera calcite	Li, B, Mg, Mn, Zn, Sr, Cd, U	Yu et al., 2007
rinsing, H ₂ O ₂ /NaOH, NH ₂ OH·HCl /NH ₃ , Na ₂ S ₂ O ₄ /citric acid, methanol	ostracod calcite	Mg, Sr	Jin et al., 2006
H ₂ O ₂ , NaOCl, heat, NH ₂ OH·HCl /NH ₃ , Na ₂ S ₂ O ₄ /citric acid	ostracod calcite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Mg, Sr	Keatings et al., 2006
hydrazine/ammonia/citrate	foraminifera calcite	Mg, Mn, Fe, Sr	Weldeab et al., 2006
rinsing, H ₂ O ₂ , NaOCl	coral aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Grottoli et al., 2005
rinsing, H ₂ O ₂ /NaOH, HNO ₃ , methanol, hydrazine/ammonia/citrate,	foraminifera calcite	Mg, Al, Mn, Sr	Pena et al., 2005
H ₂ O ₂	foraminifera calcite	$\delta^{18}\text{O}$	Waelbroeck et al., 2005
rinsing, H ₂ O ₂ /NaOH, HNO ₃	coral aragonite	Cu, Sn	Inoue et al., 2004a
HNO ₃ , ethanol, H ₂ O ₂ /NaOH/hydrazine/am	foraminifera calcite	Mg	Klinkhammer et al., 2004

monia/citrate			
rinsing, H ₂ O ₂ /NaOH, NaOCl	foraminifera calcite	Mg, Mn, Cd, Sr, Ba, U	Pak et al., 2004
rinsing, H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/citrate	foraminifera calcite	Mg, Mn, Fe, Sr	Barker et al., 2003
H ₂ O, H ₂ O ₂ , NaOH, NaOCl	coral aragonite	microscopy	Clode and Marshall, 2003
rinsing (H ₂ O + methanol)/HNO ₃ , H ₂ O ₂ /NaOH, hydrazine/ammonia/citrate, DTPA/NaOH	foraminifera calcite	Mg, Sr	Martin and Lea, 2002
rinsing, H ₂ O ₂ , HNO ₃	coral aragonite	Na, Mg, Sr	Mitsuguchi et al., 2001
rinsing, H ₂ O ₂ /NaOH, H ₂ O ₂ /NaOCl, TMAH/H ₂ O, TMAH/ethanol, natural oxidation	abiotic and coccolith calcite, organics	Mg	Stoll et al., 2001
rinsing, H ₂ O ₂ , H ₂ O ₂ /NaOH, HNO ₃ , NaOH, NaOCl	coral aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Mg, Sr	Watanabe et al., 2001
rinsing, HNO ₃ , combinations thereof	otolith aragonite	Li, Mg, Zn, Sr, Ba	Campana et al., 2000
rinsing, H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/citrate	foraminifera calcite	Mg	Hastings et al., 1998
H ₂ O ₂ , heat	coral aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Boiseau and Juillet-Leclerc, 1997
H ₂ O ₂ /(NaPO ₃) _n , HNO ₃ , NaOCl	coral aragonite	organic carbon	Hart et al., 1997
treatment order for: hydrazine/ammonia/citrate + H ₂ O ₂ /NaOH	foraminifera calcite	Cd	Rosenthal et al., 1997
H ₂ O ₂ /NaOH, hydrazine/ammonia/citrate	Halimeda aragonite	Mg, Mn, Sr, Cd	Delaney et al., 1996
H ₂ O ₂ , heat	mollusc shell powders	CO ₂ , Na, Mg, SO ₄ , Cl, Ca, Sr, crystallography	Lecuyer, 1996
rinsing, H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/citrate	coral aragonite	U	Shen and Dunbar, 1995
H ₂ O ₂ , H ₂ O ₂ /NaOH, NaOH, NaOCl	biotic calcite and aragonite	N, C, microscopy, water, organics	Gaffey and Bronnimann, 1993


DTPA, hydrazine/ammonia/citrate	foraminifera calcite, coral aragonite	Ba	Lea and Boyle, 1993
H ₂ O, H ₂ O ₂ , NaOH, NaOCl, ethanol, acetone, detergent	abiotic calcite and aragonite	Ca release, crystallography	Pingitore et al., 1993
rinsing, NH ₂ OH·HCl /NH ₃ , Na ₂ S ₂ O ₄ /citric acid	ostracod calcite	Mg, Ca, Mn, Fe, Sr	Holmes, 1992
heat (various temperatures)	biogenic carbonates (various species)	Mg, S, Ca, Sr, microscopy, crystallography, water content, hydrogen bonding, hydroxide, organics, S oxidation state	Gaffey et al., 1991
NaCl, NaHCO ₃ , Na ₂ CO ₃ , (NaPO ₃) _n , Na ₅ P ₃ O ₁₀	ostracod calcite	microscopy	Kontrovitz et al., 1991
H ₂ O ₂ , NaOH, NaOCl, heat	abiotic aragonite	Na, Mg, K, Ca, Mn, Fe, Sr	Love and Woronow, 1991
rinsing, H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/citrate	coral aragonite	Mn	Shen et al., 1991
H ₂ O ₂ /NaOH, HNO ₃ , hydrazine	coral aragonite	Cu	Linn et al., 1990
H ₂ O ₂ /methonal/heat with/without sonication	foraminifera calcite, marble, organics	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Sarkar et al., 1990
H ₂ O ₂ /NaOH, heat, grinding	pteropod and heteropod aragonite	Pb, Po	Fabry and Delaney, 1989
H ₂ O ₂ , heat	coral aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	McConnaughey, 1989
rinsing, H ₂ O ₂ /NaOH, HNO ₃ , hydrazine/ammonia/citrate	coral aragonite	Zn, Cd, Pb	Shen and Boyle, 1988
NaOCl, heat	pteropod aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Grossman et al., 1986
acetate, NH ₂ OH·HCl /NH ₃ , Na ₂ S ₂ O ₄ /citric acid, CO ₂	foraminifera calcite	Mg, Al, Ca, Mn, Fe, Ni, Cu, Zn, Cd	Boyle, 1981
heat	foraminifera	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Erez and Honjo,

	calcite		1981
H ₂ O ₂ /NH ₃ , heat	foraminifera calcite	$\delta^{13}\text{C}$	Ganssen 1981
NaOCl	ostracod calcite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Durazzi, 1977
NaOCl, heat	coral aragonite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Land et al., 1975
heat	bivalve and abiotic CaCO ₃	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Mook, 1971
NaOCl	foraminifera calcite	$\delta^{13}\text{C}$, $\delta^{18}\text{O}$	Duplessy et al., 1970
NaOCl, heat	biotic and abiotic CaCO ₃	$\delta^{18}\text{O}$	Emiliani, 1966

1164

1165

1166



1167

1168 Figure S1. Solution chemistry (pH_T , alkalinity) during growth of synthetic aragonites. In
 1169 both experiments the ^{87}Sr isotope spike was added first, ^{86}Sr second. Experiment 1/19/11
 1170 e is shown in open symbols, 1/19/11 d in filled. Symbols represent individual
 1171 measurements.

1172






Figure S2. Element ratios for a foraminifera sample prepared with different cleaning protocols. Values are mmol Sr, Mg, Na, and B, or μ mol Pb, U, Ba, and Li per mol Ca; symbols are means, error bars are 2 standard errors.




1179

1180 Figure S3. Element ratios for a *Porites sp.* sample containing tissue prepared with
 1181 different cleaning protocols. The intensive bleaching procedure is represented by
 1182 bleachx2. Values are mmol Sr, Mg, or μmol U and Ba per mol Ca; symbols are means,
 1183 error bars are 2 standard errors.


1184



1187 Figure S4. Element ratios for a synthetic aragonite sample (TS3) prepared with different
 1188 cleaning protocols (done at UWA). Values are mmol Sr, Mg, Na, and B, or μmol Pb, U,
 1189 and Ba per mol Ca; symbols are means, error bars are 2 standard errors.
 1190
 1191



1192
 1193 Figure S5. Element ratios for a synthetic aragonite sample (TS3) prepared with different
 1194 cleaning protocols (done at WHOI). Cleaning protocols consist of water rinses (DI) and
 1195 basic cleaning solutions (NaOH or NaOCl) used in different combinations and
 1196 concentrations. Values are mmol Sr, Mg, or μmol U and Ba per mol Ca; symbols are
 1197 means, error bars are 2 standard errors.
 1198



1200

1201 Figure S6. SEM image of precipitate 1/19/11 e. Note the relatively small crystal size and
1202 exposed nucleation sites.

1203

1204




Figure S7. Relative mass loss versus $\delta^{11}\text{B}$, $^{84}\text{Sr}/\text{Ca}$, $^{86}\text{Sr}/\text{Ca}$, $^{87}\text{Sr}/\text{Ca}$, Na/Ca , and Mg/Ca for synthetic aragonites. Up triangles represent 1/19/11 e (plotted on the axis to the right of each plot), down triangles 1/19/11 d (plotted on the left hand axis). Different treatments are shown in different colors. Error bars are the greater of 1 standard deviation for repeated measures of a given sample, or the measurement standard deviation for trace element measurements. For boron isotopes, error bars are 1 standard deviation of repeated measures of the given sample, or if only one measurement was made, the average standard deviation was used.

1215

1216 A supplemental data file containing data associated with each figure is available

1217 (Supplemental file `cleaningpapersupplement.xls`).

1218

1219

1220

1221